

# International Union of Pharmacology. LXX. Subtypes of $\gamma$ -Aminobutyric Acid<sub>A</sub> Receptors: Classification on the Basis of Subunit Composition, Pharmacology, and Function. Update

RICHARD W. OLSEN AND WERNER SIEGHART

*Department of Molecular and Medical Pharmacology, Geffen School of Medicine at UCLA, Los Angeles, California (R.W.O.); and Division of Biochemistry and Molecular Biology, Center for Brain Research, Medical University of Vienna, Vienna, Austria (W.S.)*

Abstract	243
I. Introduction: definition of GABA <sub>A</sub> receptors	244
A. Ionotropic GABA-gated anion channels	244
B. The Cys-loop pentameric ligand-gated ion channel superfamily	245
C. The GABA <sub>A</sub> receptor family of 19 genes	245
D. The $\rho$ subunits and the GABA <sub>C</sub> receptor concept: not recommended	245
II. Structural basis of receptor classification	246
A. GABA <sub>A</sub> receptor family and superfamily (homology of structure and function)	246
1. Subunit gene list	247
2. Subunit splice variants	247
B. Heteropentameric assembly produces complex subtype heterogeneity	247
C. Criteria for inclusion on a list of native receptor subtypes	248
1. Subtypes based on structure, pharmacology, and function: nomenclature guidelines	248
2. Discussion of the criteria	248
III. Working list of native GABA <sub>A</sub> receptor subtypes	250
A. Evidence for subtypes from localization, abundance, subunit composition, and stoichiometry data	250
B. Pharmacological evidence for subtypes	251
1. Benzodiazepine site ligands distinguish between subtypes based on $\alpha$ and $\gamma$ subunits	251
2. Evidence from other ligands (GABA, picrotoxin, general anesthetics, and ethanol) confirms and extends the list of receptors	253
C. Other, nonligand, delineation of heterogeneity	255
1. Subunit composition	255
a. Minor subunits ( $\rho$ 1-3, $\gamma$ 1, $\gamma$ 3, $\epsilon$ , $\theta$ , and $\pi$ )	255
b. $\alpha/\beta$ pentamers	256
c. Multiple $\alpha$ or $\beta$ (other?) isoforms per pentamer	256
d. Receptors containing 1, 2, 4, or 5 different subunits	256
2. Localization, trafficking, post-translational modifications, and associated proteins	256
D. Tentative list of naturally occurring receptor subtypes	257
IV. Concluding remarks	257
References	257

**Abstract**—In this review we attempt to summarize experimental evidence on the existence of defined native GABA<sub>A</sub> receptor subtypes and to produce a list of

receptors that actually seem to exist according to current knowledge. This will serve to update the most recent classification of GABA<sub>A</sub> receptors (*Pharmacol Rev* 50:291-313, 1998) approved by the Nomenclature Committee of the International Union of Pharmacology. GABA<sub>A</sub> receptors are chloride channels that mediate the major form of fast inhibitory neurotransmission in the central nervous system. They are members of the Cys-loop pentameric ligand-gated ion channel (LGIC) superfamily and share struc-

Address correspondence to: Dr. Richard W. Olsen, Department of Molecular and Medical Pharmacology, Geffen School of Medicine at UCLA, Room CHS 23-120, 650 Young Drive South, Los Angeles, CA 90095-1735. E-mail: rolsen@mednet.ucla.edu

This article is available online at <http://pharmrev.aspetjournals.org>.  
doi:10.1124/pr.108.00505.

tural and functional homology with other members of that family. GABA<sub>A</sub> receptors are assembled from a family of 19 homologous subunit gene products and form numerous, mostly hetero-oligomeric, pentamers. Such receptor subtypes with properties that depend on subunit composition vary in topography and ontogeny, in cellular and subcellular localization, in their role in brain circuits and behaviors, in their mechanisms of regulation, and in their pharmacology. We propose several criteria, which can be

applied to all the members of the LGIC superfamily, for including a receptor subtype on a list of native hetero-oligomeric subtypes. With these criteria, we develop a working GABA<sub>A</sub> receptor list, which currently includes 26 members, but will undoubtedly be modified and grow as information expands. The list is divided into three categories of native receptor subtypes: "identified," "existence with high probability," and "tentative."

## I. Introduction: Definition of GABA<sub>A</sub> Receptors

$\gamma$ -Aminobutyric acid (GABA<sup>1</sup>), the major inhibitory neurotransmitter in the brain, exerts its action via ionotropic GABA<sub>A</sub> and metabotropic GABA<sub>B</sub> receptors. GABA<sub>A</sub> receptors (GABA<sub>A</sub>-Rs) are the major inhibitory receptors in the central nervous system (CNS). They were first identified pharmacologically as being activated by GABA and the selective agonist muscimol, blocked by bicuculline and picrotoxin, and modulated by benzodiazepines, barbiturates, and certain other CNS depressants (Macdonald and Olsen, 1994; Sieghart, 1995). GABA<sub>B</sub> receptors are activated by GABA and the selective agonist baclofen but are not sensitive to typical GABA<sub>A</sub>-R ligands, such as bicuculline and muscimol. GABA<sub>B</sub> receptors couple to several different effector systems, depending on the associated G protein (e.g., G<sub>i/o</sub>), such as activation of inwardly rectifying K<sup>+</sup> channels or inhibition of high voltage-activated Ca<sup>2+</sup> channels, involving regulation of cyclic AMP or inositol phosphate signaling (Bowery et al., 2002; Bettler et al., 2004).

The objective of this review is to define GABA<sub>A</sub>-Rs and, in so doing, to summarize classifications and provide guidelines on nomenclature. This should serve to update the nomenclature suggested by Barnard et al. (1998), which remains useful and relevant, for most issues. GABA<sub>A</sub>-Rs are chloride channels that are gated by GABA and mediate rapid phasic inhibitory synaptic

transmission and also tonic inhibition by producing current in extrasynaptic and perisynaptic locations (Mody and Pearce, 2004; Farrant and Nusser, 2005). They are abundant in the nervous system of all organisms with a nervous system, including invertebrates (Buckingham et al., 2005), which also express GABA<sub>A</sub>-Rs on some muscle cells (Robinson and Olsen, 1988), where they also mediate Cl<sup>-</sup>-dependent inhibition. GABA<sub>A</sub>-Rs are found, but in a limited capacity, in non-neural tissues [such as the pancreas (Borboni et al., 1994)], where their functional roles are still under study and their pharmacological relevance remains to be established. Consistent with the emphasis of the Nomenclature Committee of the International Union of Pharmacology (IUPHAR), the receptors considered here are limited to mammalian species, with emphasis on humans. Because of their widespread localization throughout the mammalian nervous system, GABA<sub>A</sub>-Rs play a major role in virtually all brain physiological functions and serve as targets of numerous classes of drugs, used both clinically and important as research tools. The GABA<sub>A</sub>-Rs are a family of probably many receptor subtypes, but so far only a few dozen subtypes have been identified with reasonable certainty. Because of recent advances in knowledge of their molecular makeup, identification of native subunit compositions, and relevance to pharmacological specificity, an update of the list of GABA<sub>A</sub>-Rs is in order. This list is necessarily a continually changing work in progress.

<sup>1</sup> Abbreviations: GABA,  $\gamma$ -aminobutyric acid; GABA<sub>A</sub>-R, GABA type A receptor; CNS, central nervous system; IUPHAR, International Union of Pharmacology; LGIC, ligand-gated ion channel; nAChR, nicotinic acetylcholine receptor; GlyR, glycine receptor; 5-HT, serotonin; 5HT<sub>3</sub>R, 5-HT<sub>3</sub> receptor; FRET, fluorescence resonance energy transfer; RT-PCR, reverse transcriptase-polymerase chain reaction; flumazenil, ethyl-8-fluoro-5,6-dihydro-5-methyl-6-oxo-4H-imidazo[1,5-a][1,4]benzodiazepine-3-carboxylate; Ro15-4513, (3-ethyl-8-azido-5,6-dihydro-5-methyl-6-oxo-4H-imidazo[1,5-a][1,4]-benzodiazepine-3-carboxylate; BZ, benzodiazepine; CL218-872, triazolopyridazine, 3-methyl-6-[3-(trifluoromethyl)phenyl]-1,2,4-triazolo[4,3-b]pyridazine; L838,417, 7-*tert*-butyl-3-(2,5-difluorophenyl)-6-(2-methyl-2H-[1,2,4]triazol-3-ylmethoxy)-[1,2,4]triazolo[4,3-b]pyridazine; TPA003, 4,2'-difluoro-5'-[8-fluoro-7-(1-hydroxy-1-methylethyl)imidazo[1,2-a]pyridine-3-yl]biphenyl-2-carbonitrile; TPA023, 7-(1,1-dimethylethyl)-6-(2-ethyl-2H-1,2,4-triazol-3-ylmethoxy)-3-(2-fluorophenyl)-1,2,4-triazolo[4,3-b]pyridazine;  $\alpha$ 5IA, 3-(5-methylisoxazol-3-yl)-6-[(1-methyl-1,2,3-triazol-4-yl)methoxy]-1,2,4-triazolo[3,4-a]phthalazine; SR95531, gabazine, 2-(3'-carboxy-2'-propyl)-3-amino-6-*p*-methoxyphenylpyridazinium bromide; gaboxadol, 4,5,6,7-tetrahydroisoxazolo[5,4-*c*]pyridin-3-ol (THIP).

### A. Ionotropic GABA-Gated Anion Channels

Mammalian GABA<sub>A</sub>-Rs are all anion-selective channels. Increased chloride permeability generally reduces neuronal excitability (inhibition), because the Cl<sup>-</sup> equilibrium potential in most mature neurons is near the resting membrane potential and the concentration of chloride within the neuron ([Cl<sup>-</sup>]<sub>i</sub>) is much less than that within the extracellular fluid ([Cl<sup>-</sup>]<sub>o</sub>) (Martin and Olsen, 2000). However, depending on expression of Cl<sup>-</sup> transporters, [Cl<sup>-</sup>]<sub>i</sub> can increase, leading to a Cl<sup>-</sup> equilibrium potential that is less negative than the resting membrane potential. Under such conditions, activation of GABA<sub>A</sub>-Rs can cause membrane depolarization, possibly sufficient to elicit action potential discharge (excitation). The situation occurs in nature and is especially relevant in early development (Ben-Ari, 2002). In addition, on strong activation of GABA<sub>A</sub>-Rs, the resulting

increase in  $[Cl^-]_i$  might shift the membrane potential toward the firing threshold, causing rebound excitation of neurons (Marty and Llano, 2005). GABA<sub>A</sub>-R channels can conduct other anions with variable permeability ratios relative to  $Cl^-$ .  $HCO_3^-$  flux could be physiologically relevant under certain conditions (Kaila et al., 1997). The importance of bicarbonate varies with tissue and is dependent on carbonic anhydrase activity, including the tissue isozyme expression, as well as anion pumps (Rivera et al., 2005).

### B. The Cys-Loop Pentameric Ligand-Gated Ion Channel Superfamily

GABA<sub>A</sub>-Rs are part of the Cys-loop pentameric ligand-gated ion channel (LGIC) superfamily, including nicotinic acetylcholine receptors (nAChRs) (Corringer et al., 2000; Lukas and Bencherif, 2006), glycine receptors (GlyRs) (Breitinger and Becker, 2002), ionotropic 5-HT receptors (5HT<sub>3</sub>Rs) (Davies et al., 1999; Thompson and Lummis, 2006), and a  $Zn^{2+}$ -activated ion channel (Davies et al., 2003). They differ in structure from two additional LGIC families: the tetrameric glutamate receptors (P2X) (Chen and Wyllie, 2006) and the trimeric purine receptors (Khakh et al., 2001; Khakh and North, 2006). Whereas the nAChR, 5-HT<sub>3</sub>R, and the  $Zn^{2+}$ -activated channels are cation-selective channels and thus excitatory, the GABA<sub>A</sub>-R and GlyR families are anion-selective channels and, thus, with the exceptions noted above, mediate inhibition. All of the subunit members of the Cys-loop LGIC superfamily show sequence homology on the order of 30% identity but even greater similarity at the level of secondary and tertiary structure. Such receptors are all organized as pentameric membrane-spanning proteins surrounding a central pore that forms the ion channel through the membrane (Fig. 1). They all use similar sequences and functional domains to establish membrane topology, ion channel structure, agonist binding sites, and even binding sites for diverse allosteric ligands. Each subunit consists of a long N-terminal extracellular hydrophilic region, followed by four trans-membrane (M)  $\alpha$ -helices with a large intracellular loop between M3 and M4, and ends with a relatively short extracellular C-terminal domain. M2 forms the lining of the ion channel, with a possible contribution from M1 (Corringer et al., 2000; Sine and Engel, 2006). An  $\alpha$ -helical domain within the M3–M4 cytoplasmic loop has also been shown to influence ion conduction (Peters et al., 2005). The structure of the Cys-loop LGIC superfamily has been investigated by a large number of biochemical approaches, with current work emphasizing domains for ligand binding and coupling to ion channel gating, as well as subunit-subunit interactions and investigation of the role of intracellular domains. All of these data were confirmed and fell into place when combined with X-ray crystallography data on the snail soluble acetylcholine binding protein (Brejc et al., 2001), recently followed by that of the water-soluble portion of




FIG. 1. Schematic rendition of a Cys-loop pentameric ligand-gated ion channel receptor. The different shades of gray signify different subunits (unspecified here), assembled in different permutations to generate multiple hetero-oligomers. The purpose of this review is to attempt to list the naturally occurring subtypes of GABA<sub>A</sub> receptors. Similar chores face those interested in other members of the superfamily as well as other structural types of LGIC. [Adapted from Haefely W (1987) Structure and function of the benzodiazepine receptor. *Chimia* 41:389–396. Copyright © 1987 Swiss Chemical Society. Used with permission.]

the nAChR  $\alpha 1$  subunit (Dellisanti et al., 2007). The complete channel structure has been progressively well resolved (currently 4 Å) for the nAChR of *Torpedo marmorata* obtained by cryoelectron microscopy and image reconstruction (Miyazawa et al., 2003, Unwin, 2005).

### C. The GABA<sub>A</sub> Receptor Family of 19 Genes

With the complete sequence of the genome for human and a few other vertebrate species, it is now clear that there are 19 genes for GABA<sub>A</sub>-Rs (Fig. 2) (Simon et al., 2004). These include 16 subunits ( $\alpha 1$ –6,  $\beta 1$ –3,  $\gamma 1$ –3,  $\delta$ ,  $\epsilon$ ,  $\theta$ , and  $\pi$ ) combined as GABA<sub>A</sub> and 3  $\rho$  subunits, which contribute to what have sometimes been called GABA<sub>C</sub> receptors. Birds and probably some other species additionally express  $\beta 4$  and  $\gamma 4$ , but lack  $\theta$  and  $\epsilon$  subunits, so also total 19. The sequence of  $\epsilon$  is closely related to  $\gamma$  subunits (Fig. 2) and  $\theta$  to  $\beta$ , suggesting an evolutionary relationship, perhaps more evident in birds than mammals. The list is modified from Barnard et al. (1998) in that we include the  $\theta$ , but not the  $\beta 4$ , subunit.

### D. The $\rho$ Subunits and the GABA<sub>C</sub> Receptor Concept: Not Recommended

“GABA<sub>C</sub> receptors” were originally described precloning, based on a distinctive pharmacology, to encompass




FIG. 2. Dendrogram of the known 19 genes for human GABA<sub>A</sub> receptors. The distances along each line are proportional to the degree of sequence identity between the different homologous subunits. The Greek letters signify subunit families of high (>70%) identity, with different Greek letter subunit families showing homology but lower sequence identity. The distances reflect the evolutionary times required to generate sufficient sequence divergence. [Reproduced from Simon J, Wakimoto H, Fujita N, Lalonde M, and Barnard EA (2004) Analysis of the set of GABA<sub>A</sub> receptor genes in the human genome. *J Biol Chem* 279:41422–41435. Copyright © 2004. Used with permission.]

responses to GABA that did not fit either the “A” (blocked by bicuculline) or “B” (activated by baclofen) category. Such “nonA, nonB” GABA responses were found in numerous regions of the CNS. Attempts to find such a nonA, nonB GABA-activated current by expressing poly(A)<sup>+</sup> RNA from brain in oocytes were unsuccessful but a cDNA expressing this sort of GABA-evoked current was isolated from retina (Woodward et al., 1992). After cloning of the various GABA<sub>A</sub> receptor subunit genes, it was demonstrated that the ρ subunits are closely related in sequence, structure, and function to the other GABA<sub>A</sub>-R subunit families designated with other Greek letters and thus qualify for inclusion within that family. It is now clear that there are three subtype genes of ρ subunits that make homopentameric chloride channels. Such ρ-receptors show some of the pharmacological properties of the GABA<sub>C</sub> receptors, such as sensitivity to the GABA analog, CACA (*cis*-aminocrotonic acid). They are relatively insensitive to bicuculline and also to GABA<sub>A</sub>-R modulators such as benzodiazepines, barbiturates, and general anesthetics, at appropriate concentrations. GABA<sub>C</sub> receptors are sensitive to picrotoxin, neurosteroids, and some other drugs, but the overall pharmacology differs from that of most traditional

GABA<sub>A</sub>-Rs (Bormann and Feigenspan, 1995; Johnston, 1996). The ρ subunits are not simply equivalent to GABA<sub>C</sub> receptors, because some regions of the nervous system seem to lack ρ subunits and yet exhibit GABA<sub>C</sub> (i.e., non-A, non-B) pharmacology. The ρ subunits are all expressed primarily in the retina, but unlike the ρ1, the ρ2 and ρ3 subunits are also found elsewhere (Johnston, 2002). At this time it is not clear whether ρ1–3 can combine with each other in heteromers. Enz and Cutting (1999) claimed that recombinant ρ1 and ρ2 subunits can form hetero-oligomers with distinct physical properties. In native receptors this is difficult to prove because there are no antibodies that distinguish between different ρ subunits. Nor is it established whether ρ1–3 can combine in nature with members of the other 16 GABA<sub>A</sub>-R subunits. Some evidence suggests these possibilities but it is not decisive (Sieghart and Ernst, 2005). For example, coassembly of ρ and γ2 subunits was reported (Milligan et al., 2004; Pan and Qian 2005), and some cells in hippocampus show GABA<sub>A</sub>-R-like properties intermediate between ρ and γ2-containing GABA<sub>A</sub>-Rs (Hartmann et al., 2004). There is also some evidence for association of ρ with GlyRs (Pan et al., 2000).

The close structural similarities of ρ subunits to the other GABA<sub>A</sub>-R subunits, the similarities in anion channel structure and function, the important fact that other subtypes of GABA<sub>A</sub>-Rs differ in pharmacology, such as benzodiazepine sensitivity, from each other to a similar degree as do the ρ receptors, and the possibility of ρ subunits partnering with other GABA<sub>A</sub>-R subunits, led to the decision of the Nomenclature Committee of IUPHAR to designate the GABA ρ receptors as part of the GABA<sub>A</sub>-R family and to recommend against the use of the term GABA<sub>C</sub> receptor. It is especially recommended that the name GABA<sub>C</sub> receptor should not be used as the sole name for the ρ receptors in an article including, especially, the title and abstract. The ρ subunits are also discussed in section III.C.1.a.

## II. Structural Basis of Receptor Classification

### A. GABA<sub>A</sub> Receptor Family and Superfamily (Homology of Structure and Function)

The sequence homology of all 19 GABA<sub>A</sub>-R gene products to each other and to the Cys-loop pentameric LGIC superfamily lends itself to attempts to classify the receptors collectively. All members of the superfamily are homologous not only in the domains specifying membrane topology but also at the functional domain level, including ligand binding sites (Sigel and Buhr, 1997; Corringer et al., 2000; Olsen and Sawyer, 2004; Sine and Engel, 2006). Thus, homologous M2 residues are involved in channel structure and ion selectivity (Keramides et al., 2004), and the same multiple loops of extracellular sequence domains contribute to agonist/antagonist binding pockets (Galzi and Changeux, 1994; Sigel, 2002). This allows homology structural modeling

of the various functional domains comparing GABA<sub>A</sub>-Rs and nAChRs (Cromer et al., 2002; Ernst et al., 2003; Ernst et al., 2005). Even the binding sites for some allosteric modulators also show considerable homology with agonist binding sites. For example, the major benzodiazepine binding site lies at the  $\alpha/\gamma$  subunit interface and involves residues homologous to the agonist binding loops at the  $\beta/\alpha$  interface, which in turn are homologous in all members of the superfamily (Smith and Olsen, 1995; Sigel and Buhr, 1997; Corringer et al., 2000).

1. *Subunit Gene List.* The GABA<sub>A</sub>-R receptor subunit genes form a family of 19 (Table 1).

2. *Subunit Splice Variants.* Splice variants have been reported for only a few GABA<sub>A</sub>-R subunits, primarily  $\gamma 2$  (Whiting et al., 1990; Kofuji et al., 1991). The  $\gamma 2$  subunit splice variants differ in only an eight-amino acid stretch of the large intracellular loop that is present in the  $\gamma 2L$  and missing in the  $\gamma 2S$  subunit. The sequence includes a consensus protein kinase C phosphorylation substrate serine. No functional consequence of phosphorylation of the unique  $\gamma 2L$  serine residue has been convincingly demonstrated. The  $\gamma 2$  splice variants are both expressed and show differential abundance in different brain regions (Gutiérrez et al., 1994; Meier and Grantyn, 2004). In addition, they show differential aging-related changes in their level of expression (Gutiérrez et al., 1996). Brain membranes from genetically engineered mice expressing only the  $\gamma 2S$  exhibit increased affinity for benzodiazepine agonists, an effect paralleled by increased sensitivity to such compounds in behavioral responses [e.g., increased “sleep” times (loss of righting reflex)] in null allele mice (Quinlan et al., 2000). The subunit composition of the receptors responsible for such changes is unknown, and, indeed, receptors produced in compensation might include normally non-native oligomers.

Alternative splice products in the intracellular loop in  $\beta 2$  and  $\gamma 3$  subunits are found that, like  $\gamma 2L$ , also include

consensus phosphorylation substrate sequences. Alternative start sites create multiple mRNA species for  $\beta 3$ ,  $\alpha 3$ , and  $\alpha 5$  subunits, and variants lacking one or more exons have been found for  $\rho 1$ ,  $\alpha 4$ ,  $\beta 2$ , and  $\epsilon$  subunits (reviewed in Simon et al., 2004). To date, and this could change, none of these variant polypeptides have been demonstrated to be present within functional receptors, nor do they confer any unique function or pharmacology in recombinant expression systems. There is at least one report of RNA editing in the GABA<sub>A</sub>-R family (Ohlson et al., 2007), but no evidence so far for functional relevance.

### B. Heteropentameric Assembly Produces Complex Subtype Heterogeneity

The assembly of GABA<sub>A</sub>-R as heteropentamers produces complex subtype heterogeneity in structure, which is the major determinant of their pharmacological profile. These various subtypes differ in abundance in cells throughout the nervous system and thus in functions related to the circuits involved. A major factor in producing heterogeneity is the existence of the six different  $\alpha$  subunit variants. Some of these physiological receptor subtypes containing specific  $\alpha$  subunits can be distinguished by ligands that act at the benzodiazepine site (Barnard et al., 1998). In addition, there are now clear examples of  $\beta$  and  $\gamma$  or  $\delta$  or other subunit selectivity for drug action on GABA<sub>A</sub>-R subtypes (see section III.B.2). Clearly it is the nature, stoichiometry, and arrangement of the subunits that determine details of pharmacological selectivity. Thus, pharmacology can often provide evidence for structural heterogeneity and informs us about the subunit composition of native receptor subtypes.

Evidence is greatly in favor of a pentameric receptor and most GABA<sub>A</sub>-R subtypes are formed from two copies of a single  $\alpha$ , two copies of a single  $\beta$ , and one copy of another subunit, such as  $\gamma$ ,  $\delta$ , or  $\epsilon$  (Sieghart and Sperk, 2002; Olsen and Sawyer, 2004). At this time the subunit composition, stoichiometry, and wheel alignment are not known for most pharmacological subtypes: tentative definitions of subtypes can be provided as a work in progress. Current knowledge allows elimination of most of the thousands of permutations theoretically possible for combinations of the known 19 subunits into five-part (pentameric) complexes. Barnard et al. (1998) suggested a maximum on the order of 800 combinations and probably far fewer in reality on the basis of current knowledge of identified subunit partnering and apparent rules of assembly in neurons. Very few combinations have been conclusively identified *in situ*. We suggest some criteria for inclusion in a list of native subtypes and begin to generate such a list. Current evidence suggests that only 11 subtypes can be listed as conclusively identified, and these are reasonably abundant. We have also listed several subtypes for which the evidence is strong but not conclusive (six subtypes). We finish by mentioning subtypes containing one of each of the minor sub-

TABLE 1  
GABA<sub>A</sub> receptor subunit gene list

Chromosome data from Simon et al. (2004).

Subunit	Gene	Chromosome		
		Human	Mouse	Rat
$\alpha 1$	<i>GABRA1</i>	5q34	11	10
$\alpha 2$	<i>GABRA2</i>	4p12	5	14
$\alpha 3$	<i>GABRA3</i>	Xq28	X	X
$\alpha 4$	<i>GABRA4</i>	4p12	5	14
$\alpha 5$	<i>GABRA5</i>	15q13.2	7	1
$\alpha 6$	<i>GABRA6</i>	5q34	11	10
$\beta 1$	<i>GABRB1</i>	4p12	5	14
$\beta 2$	<i>GABRB2</i>	5q34	11	10
$\beta 3$	<i>GABRB3</i>	15q13.2	7	1
$\gamma 1$	<i>GABRG1</i>	4p12	5	14
$\gamma 2$	<i>GABRG2</i>	5q34	11	10
$\gamma 3$	<i>GABRG3</i>	15q13.2	7	1
$\delta$	<i>GABRD</i>	1p36.3	4	5
$\epsilon$	<i>GABRE</i>	Xq28	X	X
$\theta$	<i>GABRQ</i>	Xq28	X	X
$\pi$	<i>GABRP</i>	5q35.1	11	10
$\rho 1$	<i>GABRR1</i>	6q15	4	5
$\rho 2$	<i>GABRR2</i>	6q15	4	5
$\rho 3$	<i>GABRR3</i>	3q12.1	16	11

units, evidence for whose native existence is tentative (another eight, plus one subtype with two kinds of  $\alpha$  subunit), for a grand total of 26. An additional similar number of subtypes, which are relatively rare, but are likely to exist, are not listed at this time, but the list will continue to grow as more information becomes available. Each subtype could play a significant role in the cells in which they occur. It should be noted that even these minor GABA<sub>A</sub>-R subtypes are present in amounts comparable with or greater than receptor subtypes for accepted brain neurotransmitters other than glutamate, that is, the biogenic amines and acetylcholine. The heteromeric LGICs clearly offer much greater heterogeneity than other known receptor subtypes.

### C. Criteria for Inclusion on a List of Native Receptor Subtypes

1. *Subtypes Based on Structure, Pharmacology, and Function: Nomenclature Guidelines.* Criteria are needed to define which receptor subtypes can be accepted as being native to neurons. We suggest basing the criteria on structure, pharmacology, and function. The list of subtypes will necessarily be a work in progress as information on all LGIC families is currently incomplete. We propose five major criteria: two for recombinant studies and three for native studies, each with subclassifications, for inclusion of a subunit combination on the native receptor subtype list (Table 2). None of these criteria by itself is sufficient. Because five criteria are rarely, if ever, met, it is the remit of a committee of experts (e.g., Nomenclature Committee of the IUPHAR subcommittee) to decide which candidates qualify. At this time we choose to be relatively strict regarding inclusion, including subtypes that *either must, or are highly likely* to, exist, rather than including all subtypes that *might* exist. We have decided to include more possibilities in the list for GABA<sub>A</sub>-Rs (Table 3) by dividing it into three categories, depending on the number of criteria that are met: A) "identified," B) "existence with high probability," and C) "tentative."

2. *Discussion of the Criteria.* A brief discussion of the available pertinent evidence and its value in determining the existence of native subtypes is in order. The expression of recombinant receptors, the determination of their subunit composition and arrangement, and their biophysical and pharmacological characterization is obviously an important part in the identification of native receptors. But expression of recombinant receptors is insufficient to prove their existence *in vivo*, because gross overexpression of subunit proteins can result in heterooligomeric combinations that are unlikely to occur in nature. For example, recombinant receptors containing both the  $\gamma 2$  and  $\delta$  subunits have been generated and characterized (Saxena and Macdonald, 1994; Hevers et al., 2000), whereas the actual existence of such receptors in the brain is highly questionable (for discussion, see Sieghart and Sperk, 2002). In addition, coexpression of subunits does not always result in the exclusive forma-

TABLE 2  
Criteria for inclusion on a list of native receptor subtypes

I. Recombinant receptors
A. Evidence for their formation, subunit composition, and stoichiometry, using recombinant receptor expression in heterologous cell systems
1. The subunit polypeptides must be shown to be expressed
2. The subunit polypeptides must be shown to coassemble
a. Coimmunoprecipitation
b. Physical demonstration of subunit interactions by FRET or similar technique
c. Formation of pentamers
d. Unique subunit arrangement, e.g., using concatemers
3. The corresponding recombinant receptor subtype must be functional
B. Evidence for unique properties, including pharmacology
1. Unique biophysical characteristics
2. Unique pharmacology
a. Receptor subtype-selective agonists, antagonists, allosteric modulators
b. Receptor subtype-selective radioligands
c. Potency and efficacy for a series of ligands
d. Macrokinetic measures (e.g., apparent EC <sub>50</sub> values and binding constants for a series of ligands)
II. Native receptors
A. Colocalization of subunits
1. Tissue colocalization
2. Cell colocalization (in situ, single-cell RT-PCR)
3. Subcellular colocalization (light and electron microscopy)
B. Physical demonstration of subunit interactions (e.g., by coimmunoprecipitation)
C. Functional demonstration:
1. Evidence that a given receptor is expressed in real neurons by showing properties (assessed with electrophysiology) corresponding with a recombinant receptor candidate; microscopy may complement
2. Evidence that a given subunit or subunit combination participates in a specific function <i>in vitro</i> or <i>in vivo</i> using genetically modified mice

tion of the expected receptors. Thus, several reports have shown that coexpression of  $\alpha$ ,  $\beta$ , and other ( $\gamma$ ,  $\delta$ , and  $\epsilon$ ) subunits in oocytes or cell lines can lead to considerable expression of  $\alpha\beta$  receptor channels, without  $\gamma$ ,  $\delta$ , or  $\epsilon$ . One must either precipitate the receptors via the third component subunit measuring subunit composition by Western blots and pharmacological properties via binding studies or use a pharmacological test to demonstrate the expression of this subunit and its contribution to the channel currents. In any case, it has to be kept in mind that receptor heterogeneity might influence results of the subsequent characterization of recombinant receptors. To increase the formation of receptors containing  $\alpha$  and  $\beta$ , as well as  $\gamma$ ,  $\delta$ , or  $\epsilon$  subunits, higher amounts of mRNA or cDNA for the third component subunit often are used, as well as longer times for expression (Boileau et al., 2002, 2005; Olsen et al., 2007), despite the risk of non-natural or pathological pentamers with too many copies of a given subunit by the use of such extremes.

Thus, in recombinant expression studies, one must demonstrate that the subunits under study are indeed ex-

pressed, combine into receptors, form pentamers with defined subunit stoichiometry and arrangement, and have function. If possible, the properties of receptors should be compared with those of receptors with defined subunit composition and arrangement using concatemers. One might add that researchers also need to pay attention to the source of their receptor subunit clones and specify the source and sequences of those they use. A single point mutation in the gene of a subunit isolated from the brain could change the properties of the resulting recombinant receptors (Sigel et al., 1992). Then one can examine the receptor for unique biophysical properties using electrophysiology, assessing agonist and modulator mechanisms, determined, for example, from single channel and macroscopic current kinetics, single channel conductance, and possibly conductance substates. Finally, one can examine the receptor for unique pharmacology, using subtype-selective ligands, including radioligands, and measuring relative affinities, potencies, and efficacies for a series of ligands.

Expression in the brain at the mRNA (in situ hybridization, RT-PCR, single cell RT-PCR) or protein level (immunostaining) is a good starting point to indicate that two given subunits may be coexpressed in a given cell and which types of cells express them. A combination of in situ hybridization and immunohistochemistry is stronger than either technique alone. But colocalization of two subunits, although necessary, is not sufficient to establish whether these subunits are partners in a pentameric receptor subtype. Individual subunits alone most often do not define a receptor and might even have functions different from those of receptors. Microscopic colocalization of subunit proteins is one of the factors used in deciding native subtype composition, but colocalization at the light microscope level is not conclusive evidence for coexpression within a pentamer. Likewise, colocalization at the electron microscope level is useful evidence but does not necessarily define receptor subtypes, because even subunits located side-by-side with immunogold labeling could belong to adjacent receptors. Anatomy compendia are being developed by experts, and these play a role in deciding on inclusion of native subtypes. The questionable specificity of many antibodies, however, suggests that great care must be taken to analyze the published work on subunit localization (Rhodes and Trimmer, 2006; Moser et al., 2007). Because most antibodies used are polyclonal and because the composition of polyclonal antibodies is different in each donating animal and even in each blood sample, such verification of the specificity of the antibodies has to go on until highly selective and well characterized monoclonal or recombinant antibodies are available. All antibodies used for these compendia should thus be used in wild-type and knockout mice to be sure that they unequivocally identify the correct protein only. But even if the data are correct, most data on

localization are not in themselves sufficient for defining receptor subtype composition.

Coimmunoprecipitation of subunits comes close to defining a subtype because it indicates (but not necessarily proves) assembly. Here again, the antibodies used must be demonstrated to have total subunit specificity as any cross-reactivity contaminates the results. The antibodies must be characterized by Western blotting on crude brain tissue or cells and be shown to recognize the band of correct size, and only that, in normal but not knockout mice, when such mice are available. Specific antibody reagents are currently available for such an approach, and the provider of such antibodies should be asked for data documenting the absence of cross-reactivity. Nevertheless, incompletely assembled intermediates could contribute, and subunits or receptors could associate with each other (natural interaction, artifactual aggregation, or association via cytoskeleton proteins) without being in the same receptors, weakening the strength of the evidence based on the coimmunoprecipitation method. One needs to be careful with the choice of detergent and solubilization conditions to minimize nonspecific protein interactions without destroying oligomers. If one could demonstrate that the associated subunits were present in a detergent-solubilized protein of the correct size for a pentamer this would strengthen the argument, but this is rarely, if ever, provided, and even such studies may be ambiguous because of comigration of multiple heterooligomeric protein species, so the results in any case must be regarded with caution. In the case of G protein-coupled receptor heteromultimer possibilities, evidence for true subunit-subunit association considers physical techniques such as FRET (Pin et al., 2007), and this approach may also find use for ligand-gated ion channel receptors. However, the method is extremely difficult to apply to native proteins, requiring the use of transgenic mice. If it is performed with recombinant receptors in a heterologous expression system, there is again the possibility of identifying non-native combinations owing to subunit overexpression. But such studies again would not provide a clear-cut answer because subunits demonstrated to be close together by FRET techniques could have assembled in the same receptor or be located in two different receptors associated with each other. The above considerations seem trivial, but they are not, judged by proposals of acknowledged scientists on how to unequivocally solve the problems of establishing receptor subunit composition that do not survive more thorough discussions.

Thus, the conclusion is that a combination of different techniques has to be used to identify a receptor subunit composition of a native receptor, knowing the limits of each technique. Even then, one cannot be absolutely sure, but at least this is at the limit of the techniques available. What is really needed is "in situ" electrophysiology with a channel characterization and a pharmacological fingerprint of the receptors to prove that the

subunits known to exist in that cell contribute to the receptor studied and its physiology. This is extremely difficult, and fingerprints for the different receptor subtypes are not always sufficiently selective, may be host cell-specific, or are lacking altogether. One must take care to be as certain as possible that the subunits meant to be expressed in a recombinant system for obtaining a fingerprint are actually present and contributing to the data and that the pentamer has correct stoichiometry (see discussion above). Recombinant expression suffers from uncertainty that perhaps other gene products present in neurons might not be present in the cells used for recombinant expression (e.g., associated proteins, proteins affecting trafficking, or post-translational modifications) or that they do not exactly match that in the native system because of host cell-specific factors (Birnie and Korpi, 2007). Even a biophysical characteristic as fundamental as single-channel conductance can be affected by the expression system (Lewis et al., 1997), and anecdotes abound regarding different results for the "same receptors" expressed in two kinds of cells. Concatenated subunits may be of use in determining specific subunit composition and wheel arrangement (Im et al., 1995; Baumann et al., 2002; Minier and Sigel, 2004; Boileau et al., 2005; Sigel et al., 2006), although this approach also has a number of problems (Ericksen and Boileau, 2007).

To summarize the strategy used: one first notes which subunits are expressed in a given cell type. Then one looks at evidence that a given pair or triplet of subunits on that list are associated, based on coimmunoprecipitation, coregulation in cells, or knockout animals. The coexpression of  $\alpha 1\beta 2\gamma 2$  suggests the possibility of coordinate regulation of expression of chromosomal clusters, but the exceptions are too abundant to consider this a factor in partnering. Then one determines by recombinant expression and electrophysiology the channel kinetics and pharmacology of those subunit combinations, keeping in mind the numerous caveats listed above. Because most of these caveats have not been taken into account previously, one clearly cannot believe everything published on the properties of recombinant receptors. Native receptors with the same unique properties are then sought in the neurons recorded from brain slices, dissociated single cells, or at least cells in culture.

Genetically engineered mice (knockout and knockin) provide some evidence indicating that certain oligomeric receptors really do exist because one can correlate the loss of certain receptor responses and behaviors with the receptor subtypes addressed (Jones et al., 1997; Rudolph et al., 1999; McKernan et al., 2000; Sur et al., 2001; Vicini et al., 2001). The function of deleted genes is sometimes obvious, whereas in other cases, subtle changes in behavior need to be evaluated properly. Evidence from global knockout mice is often complicated by possible compensation, causing the observer to miss important functions. For example, deletion of the  $\alpha 1$  sub-

unit of the GABA<sub>A</sub>-R results in loss of nearly 50% of the total GABA<sub>A</sub>-R population in mouse brain and ablates fast synaptic transmission mediated by the abundant  $\alpha 1$  subunit-containing receptors, yet the null mutant mice display a phenotype that is grossly normal (Sur et al., 2001). Diverse transcriptional responses may act to abrogate the effect of the knockout and preserve neuronal excitability and network behavior (Ponomarev et al., 2006). Point mutated knockin mice with altered behavior/pharmacology are more convincing, and several important examples are included in the GABA<sub>A</sub>-R field. For an overview and discussion, see Olsen and Homanics (2000), Rudolph and Möhler (2004), Sieghart and Ernst (2005), and Atack (2005).

### III. Working List of Native GABA<sub>A</sub> Receptor Subtypes

#### A. Evidence for Subtypes from Localization, Abundance, Subunit Composition, and Stoichiometry Data

It was found early on in recombinant GABA<sub>A</sub>-R studies that robust GABA-activated channel formation occurred with combinations of  $\alpha$  and  $\beta$  subunits and also with  $\alpha$ ,  $\beta$ , and  $\gamma$  subunits. The latter turned out to be the prevalent native combination (Barnard et al., 1998). The vast majority of GABA<sub>A</sub>-Rs in the CNS contain the  $\gamma 2$  subunit, and this is the most abundant subunit in rat brain and in most regions based on *in situ* hybridization of mRNA (Wisden et al., 1992; Laurie et al., 1992; Persohn et al., 1991, 1992) and immunostaining (Fritschy and Möhler, 1995; Pirker et al., 2000). The  $\gamma 1$  and  $\gamma 3$  subunits are rarer but have some role in discrete regions, probably with well prescribed subunit partners. Thus, approximately 75 to 80% of GABA<sub>A</sub>-Rs contain the  $\gamma 2$  subunit (Whiting et al., 2000; Sieghart and Ernst, 2005).

The  $\alpha 1$  is the most abundant of the  $\alpha$  subunits and is often colocalized with its chromosome partners, the likewise highly expressed  $\beta 2$  and  $\gamma 2$  subunits (Sieghart and Sperk, 2002). As noted above, knockout of the  $\alpha 1$  subunit causes total GABA<sub>A</sub>-R content in mouse brain to decrease by 50% (Sur et al., 2001). The  $\alpha 2$  and  $\alpha 3$  subunits are moderately abundant and  $\alpha 5$  is relatively rare except in the hippocampus, as indicated by regional distribution and immunoprecipitation studies (Pirker et al., 2000; Sieghart and Sperk, 2002). The  $\alpha 4$  and  $\alpha 6$  subunits are reasonably highly expressed in forebrain and cerebellum, respectively. Among the  $\beta$  subunits,  $\beta 1$  is least common,  $\beta 2$  is most abundant and most widespread (knockout results in a 50% reduction in GABA<sub>A</sub>-Rs in mouse brain) (Sur et al., 2001), and  $\beta 3$  is reasonably highly expressed, but more discrete. Furthermore, it is more dense perinatally than in adult brain (Zhang et al., 1991; Laurie et al., 1992). The identity of the  $\beta$  subunit often cannot be determined, because on precipitation of GABA<sub>A</sub>-Rs with  $\alpha$ ,  $\gamma$ , or  $\delta$  subunit-specific antibodies in



most cases all three  $\beta$  subunits are coprecipitated. Only in some rare areas where there are cells with only one type of  $\beta$  subunit contributing to functional GABA<sub>A</sub>-Rs (Persohn et al., 1991, 1992; Wisden et al., 1992; Laurie et al., 1992; Pirker et al., 2000) can the type of  $\beta$  subunit be predicted. In these areas, however, no subsequent immunoprecipitation or electrophysiological studies have been performed to investigate possible assembly partners, and, thus, actual assembly of defined receptors and their composition cannot be derived with certainty. For this reason, it has been concluded that all of the  $\beta$  subunits exist in functional receptors, usually with only one type per pentamer (Whiting et al., 2000).

Based on the frequent colocalization in the brain (Fritschy and Möhler, 1995; Pirker et al., 2000) and in neurons (Klausberger et al., 2002), on results from coimmunoprecipitation experiments (Jechlinger et al., 1998; Pörtl et al., 2003), and on the pharmacology described in section III.B, the  $\alpha 1\beta 2\gamma 2$  subunit combination is considered to exist in the brain, probably in large amounts (Benke et al., 1991; Somogyi et al., 1996; Whiting et al., 2000). Recombinant  $\alpha 1\beta 2\gamma 2$  receptors have been shown to have a  $2\alpha\text{-}2\beta\text{-}1\gamma 2$  stoichiometry [denoted  $(\alpha 1)_2(\beta 2)_2\gamma 2$ ], and this stoichiometry is supported by coimmunoprecipitation results from rat or mouse brain (Sieghart and Sperk, 2002). It is generally assumed, without conclusive proof, that other  $\alpha$  and  $\beta$  subunits also combine with  $\gamma 2$  in combinations of  $2\alpha\text{-}2\beta\text{-}1\gamma 2$  (Sieghart and Ernst, 2005). Other  $\alpha$  subunits are also colocalized with and have been shown to coprecipitate with all  $\beta$  and the  $\gamma 2$  subunits (Sieghart and Sperk, 2002). Receptors containing these subunits also have a specific pharmacology (see section III.B) and thus also seem to exist in the brain.

The other "minor" subunits, notably  $\delta$ , are thought to be able to replace  $\gamma$  in the pentamer, as in the  $2\alpha\text{-}2\beta\text{-}\delta$  combination (Barrera et al., 2008). However,  $\alpha 4$  and  $\alpha 6$  subunits are just as often combined with the  $\delta$  subunit as with  $\gamma 2$  (Sieghart and Sperk, 2002). The  $\delta$  subunit is obligatorily partnered with the  $\alpha 6$  subunit in cerebellar granule cells and is primarily associated with the  $\alpha 4$  subunit in forebrain areas including dentate gyrus, neostriatum, some layers of cortex, and a few other areas. The  $\delta$  subunit seems to have a perisynaptic/extrasynaptic localization (Nusser et al., 1998; Peng et al., 2002; Wei et al., 2003). The  $\varepsilon$  subunit is rare but can substitute for  $\gamma$  or  $\delta$  in some areas, such as hypothalamus, and the  $\theta$  and  $\pi$  subunits are only sketchily characterized (Korpi et al., 2002; Sieghart and Sperk, 2002).

The  $\gamma 2$  subunit is required for synaptic localization of GABA<sub>A</sub>-Rs, usually associated with the  $\alpha 1$ ,  $\alpha 2$ , or  $\alpha 3$  subunits. Nevertheless, substantial numbers, possibly a majority, of  $\gamma 2$  subunit-containing GABA<sub>A</sub>-Rs are extrasynaptic, because of the much greater area of extrasynaptic membranes. This includes some of the  $\alpha 1/\alpha 2/\alpha 3$  and the great majority of  $\alpha 4/\alpha 5/\alpha 6$  combinations. In contrast, combinations in which other subunits, such as

$\delta$  and  $\varepsilon$ , replace  $\gamma 2$  are considered to be exclusively nonsynaptic. The physiological and pharmacological importance of perisynaptically and extrasynaptically localized GABA<sub>A</sub>-Rs has recently become increasingly appreciated (Mody and Pearce, 2004; Semyanov et al., 2004; Farrant and Nusser, 2005).

## B. Pharmacological Evidence for Subtypes

*1. Benzodiazepine Site Ligands Distinguish between Subtypes Based on  $\alpha$  and  $\gamma$  Subunits.* The typical GABA<sub>A</sub>-R is positively modulated by diazepam-like benzodiazepines and binds radioligands for the GABA site (e.g., [<sup>3</sup>H]muscimol), the benzodiazepine site (e.g., [<sup>3</sup>H]flunitrazepam, [<sup>3</sup>H]flumazenil, or [<sup>3</sup>H]Ro15-4513, and the picrotoxin/convulsant/ channel sites (e.g., [<sup>35</sup>S]t-butyl bicyclophosphorothionate) (reviewed by Macdonald and Olsen, 1994; Sieghart, 1995; Barnard et al., 1998; Korpi et al., 2002; Johnston, 2005). Barnard et al. (1998) listed approximately 20 chemical classes of ligand that bind to the benzodiazepine (BZ) site on GABA<sub>A</sub>-R, including the structures of approximately 40 compounds. Additional chemical classes of ligands are listed in Gardner et al. (1993), Olsen and Gordey (2000), Sieghart and Ernst (2005), Atack (2005), and Whiting (2006), which should be consulted for chemical names and structures. In each of these chemical classes compounds are available that enhance (positive allosteric modulators or BZ site agonists) or reduce (negative allosteric modulators or inverse BZ site agonists) GABA-induced chloride ion flux via the same BZ binding site. In addition, in each of these chemical classes there are compounds that do not modulate GABA-induced chloride flux, although they interact with the BZ binding site (neutral BZ site ligands or BZ site antagonists). The latter compounds, however, are able to inhibit the action of BZ site agonists or inverse agonists. In between the full BZ site agonists or inverse agonists, there are compounds that elicit less drastic allosteric enhancement or reduction of GABA-induced chloride currents; these compounds are called partial agonists or partial inverse agonists at the BZ site, respectively.

The BZ binding site is located at the interface of an  $\alpha$  and a  $\gamma$  subunit. Its pharmacology is thus influenced by both of these subunits, whereas  $\beta$  subunits, although needed to construct a channel, do not greatly affect the sensitivity of the GABA<sub>A</sub>-R to BZ site ligands (Hadingham et al., 1993). The traditional BZ site agonists (GABA-enhancing CNS depressants such as diazepam) are active on the GABA<sub>A</sub>-Rs containing a  $\gamma 2$  subunit (Pritchett et al., 1989), a  $\beta$  subunit, and one of the  $\alpha$  subunits,  $\alpha 1$ ,  $2$ ,  $3$ , or  $5$ . Receptors containing the  $\gamma 2$  subunit exhibit a higher BZ sensitivity than those containing the  $\gamma 1$  subunit (Sieghart, 1995; Khom et al., 2006); the  $\gamma 3$ -containing GABA<sub>A</sub>-Rs are modulated by some BZ ligands but with altered selectivity from those incorporating the  $\gamma 2$  subunit (Sieghart, 1995; Hevers and Lüddens, 1998). The BZ-sensitive GABA<sub>A</sub>-Rs can

be further subdivided, in that receptors containing the  $\alpha 1$  subunit have a higher sensitivity to a subpopulation of BZ site ligands, the benzodiazepines quazepam and cinolazepam (Sieghart, 1989) or nonbenzodiazepines such as zolpidem (an imidazopyridine) and a few others, including CL218-872 (triazolopyridazine), zaleplon, and indiplon, and abecarnil ( $\beta$ -carboline), (Olsen and Gordey, 2000; Korpi et al., 2002; Sieghart and Ernst, 2005). Furthermore, receptors containing the  $\alpha 2$  or  $\alpha 3$  subunit have an intermediate affinity for zolpidem, whereas those containing  $\alpha 5$  have very low affinity for this drug. The differential zolpidem affinity demonstrated by recombinant GABA<sub>A</sub>-Rs containing different  $\alpha$  subunits can also be found in the brain (Itier et al., 1996; Whiting et al., 2000; Sieghart and Sperk, 2002) and individual cells can be shown to exhibit more than one GABA<sub>A</sub>-R with varying affinity for zolpidem, depending on  $\alpha$  subunit subtype expression (Criswell et al., 1997). Again, consult the cited reviews for chemical structures and analogs.

Receptors containing the  $\alpha 4$  or  $\alpha 6$  subunits, together with  $\beta$  and  $\gamma 2$ , do not bind the traditional BZ agonists, including zolpidem, but demonstrate high affinity for some ligands, notably the imidazobenzodiazepines such as flumazenil and Ro15-4513, or bretazenil (Korpi et al., 2002). Both the potency and efficacy for BZ ligands depend on the nature of the  $\alpha$  subunit.

The benzodiazepine site ligands so far available do not distinguish well between the  $\alpha 2$  and  $\alpha 3$  or between the  $\alpha 4$  and  $\alpha 6$  subunits. All four, however, can produce functional channels *in vitro* when coexpressed with other subunits, and their differential distributions in the brain suggest they will modulate different behavioral circuitry. Behavioral effects of drugs predicted from subtype selectivity *in vitro* has been successful in a few cases, such as the  $\alpha 2/3$ -selective triazolopyridazine anxiolytics based on CL218-872: L838,417, TPA003, or TPA023 (Atack, 2005; Morris et al., 2006; Atack et al., 2006).

Subtype selectivity for drugs *in vivo* has been further confirmed in subunit-specific genetically engineered mice (Rudolph and Möhler, 2004; Atack, 2005; Whiting, 2006). Thus, based on the evidence that most of the actions of diazepam are mediated via receptors composed of  $\alpha 1\beta\gamma 2$ ,  $\alpha 2\beta\gamma 2$ ,  $\alpha 3\beta\gamma 2$ , and  $\alpha 5\beta\gamma 2$  subunits, a point mutation involving a histidine to arginine substitution was introduced into the genes of the individual  $\alpha$  subunits, rendering the respective receptors insensitive to allosteric modulation by diazepam. A comparison of drug-induced behavioral responses in the mutated and wild-type mice then allowed the identification of diazepam effects that were missing, or reduced, in the mutant mice. With this approach, it was demonstrated that  $\alpha 1\beta\gamma 2$  receptors mediate the sedative, anterograde amnesic and in part the anticonvulsant actions of diazepam (Rudolph et al., 1999; McKernan et al., 2000). These two independent studies showed exactly the same be-

havioral responses, provided the mice were tested under the same conditions (Crestani et al., 2000). The anxiolytic activity of diazepam is mediated by GABA<sub>A</sub>-Rs composed of  $\alpha 2\beta\gamma 2$  subunits (Low et al., 2000), and, under conditions of high receptor occupancy, also by  $\alpha 3$  GABA<sub>A</sub>-Rs (Dias et al., 2005; Yee et al., 2005; discussed by Möhler, 2007). The  $\alpha 3$ -selective drug TP003 also implicates a role for  $\alpha 3\beta\gamma 2$  receptors in anxiolytic (Dias et al., 2005) and anticonvulsant action (Fradley et al., 2007). The  $\alpha 2\beta\gamma 2$  and  $\alpha 3\beta\gamma 2$  receptors are also implicated in some of the muscle relaxant activities of diazepam (Low et al., 2000) and the  $\alpha 3\beta\gamma 2$  in the antiabsence effects of clonazepam (Sohal et al., 2003). The  $\alpha 3$  global knockout mice displayed a hyperdopaminergic phenotype relevant for GABAergic control of psychotic-like symptoms (Yee et al., 2005). The  $\alpha 5\beta\gamma 2$  receptors seem to influence learning and memory, shown by improved spatial memory in mice with knockout of  $\alpha 5$  subunits (Collinson et al., 2000), and trace fear conditioning was facilitated in the point-mutated  $\alpha 5$  knockin which unexpectedly showed major  $\alpha 5$  subunit knockdown in the CA1 region (Crestani et al., 2002; Rudolph and Möhler, 2004). Furthermore, the  $\alpha 5$ -selective inverse agonists such as  $\alpha 5$ IA can improve cognitive function (Atack, 2005; Dawson et al., 2006). Mice lacking the  $\alpha 5$  subunit were shown to exhibit reduced amnesic response to the intravenous general anesthetic etomidate but not to the immobilizing activity of the drug (Cheng et al., 2006). Knockout mice have also implicated several GABA<sub>A</sub>-R subtypes in the reinforcing effects of ethanol, including the  $\alpha 1$  (Boehm et al., 2004), the  $\alpha 5$  (Boehm et al., 2004; Stephens et al., 2006), the  $\delta$  (Mihalek et al., 2001), the  $\alpha 4$  (Liang et al., 2008), and the  $\alpha 6$  subunit (Hanchar et al., 2005). Years of study on  $\alpha$  subunit-dependent subtypes have led to drug candidates for modifying selective behaviors and clinical indications (Atack, 2005; Sieghart and Ernst, 2005; Whiting, 2006), and the genetically modified mice have supported this subtype selectivity. Possibly even more impressive than the drug candidates has been the clues generated regarding behavior involving specific receptor subtypes and thus specific brain circuitries (for review, see Rudolph and Möhler, 2004). Although this attribution of the varied behavioral actions of diazepam site ligands and ethanol to different receptor subtypes might only be a first approximation, and the evaluation may be somewhat tentative, it clearly indicates that receptors containing the respective subunits are present in the brain and exhibit distinct functions in different neuronal circuits, verified by consistent results with subtype-selective drugs.

In contrast to  $\alpha 1\beta\gamma 2$ ,  $\alpha 2\beta\gamma 2$ ,  $\alpha 3\beta\gamma 2$ , and  $\alpha 5\beta\gamma 2$  receptors,  $\alpha 4\beta\gamma 2$  and  $\alpha 6\beta\gamma 2$  receptors are diazepam-insensitive but are still able to bind the imidazobenzodiazepines Ro15-4513, and flumazenil, which act as BZ site inverse agonist and antagonist, respectively. Thus, these receptors can be identified by ligand-binding studies and autoradiography, using [<sup>3</sup>H]Ro15-4513 in the

absence or presence of high concentrations of diazepam or flumazenil, to mask receptors containing the  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ , or  $\alpha 5$  subunits. Such studies have clearly identified diazepam-insensitive, but flumazenil-sensitive, binding sites in cerebellum and forebrain. These sites disappear in  $\alpha 6$  or  $\alpha 4$  null mutant mice, indicating that they represent  $\alpha 6\beta\gamma 2$ , or  $\alpha 4\beta\gamma 2$  receptors, respectively (Korpi et al., 2002). In addition, there are GABA<sub>A</sub>R-mediated inhibitory currents recorded in hippocampus that are enhanced by Ro15–4513, an effect specific for  $\alpha 4\beta\gamma 2$  (Liang et al., 2006). So, even with limited pharmacology, some in vivo and in vitro information about selectivity, and the help of genetically modified mice, we can make receptor assignments; e.g., each  $\alpha$  subunit defines at least one unique subtype, partnered with a  $\beta$  and a  $\gamma 2$  subunit. Thus, GABA<sub>A</sub>-Rs exist with binding properties consistent with high, intermediate, and low affinity for certain BZ site ligands, such as zolpidem, and GABA<sub>A</sub>-R channels with this pharmacological specificity are localized in certain brain regions where the different subunits are expressed. In addition, all  $\alpha$  subunits can be coprecipitated with  $\beta$  and  $\gamma 2$  subunits in these regions. Therefore, we conclude that the six different  $\alpha$  subunits occur in the brain, combined with  $\beta$  and  $\gamma 2$ , and can start our receptor list with all six  $\alpha$  subunits, each combined with  $\beta\gamma 2$ , for a total of six subtypes (Macdonald and Olsen, 1994; Sieghart, 1995; Barnard et al., 1998; Whiting et al., 2000; Korpi et al., 2002). As discussed in section III.A, only for the  $\alpha 1\beta\gamma 2$  receptor subtype does sufficient evidence identify the type of associated  $\beta$  subunit ( $\alpha 1\beta 2\gamma 2$ ).

The  $\delta$  subunit, generally partnered with the  $\alpha 4$  and  $\alpha 6$  subunits, produces the seventh and eighth subtypes. This is supported by evidence of colocalization and coimmunoprecipitation (Jechlinger et al., 1998; Whiting et al., 2000; Pörtl et al., 2003; Sieghart and Ernst, 2005), accompanied by electrophysiological evidence for  $\alpha 6\beta\delta$  GABA<sub>A</sub>-Rs in cerebellar granule cells and  $\alpha 4\beta\delta$  in forebrain, based on tonic BZ-insensitive GABA<sub>A</sub>-R currents, which are lost in the  $\delta$  knockout mice (Stell et al., 2003; Wei et al., 2003; Belelli et al., 2005; Hanchar et al., 2005; Liang et al., 2006; Glykys et al., 2007). Such GABA<sub>A</sub>-Rs bind even fewer BZ site ligands. Nevertheless, these subtypes have recently been shown to bind Ro15–4513, flumazenil, and certain  $\beta$ -carbolines (Hanchar et al., 2006; Wallner et al., 2006). Thus, current evidence supports the existence of  $\alpha 4\beta\delta$  and  $\alpha 6\beta\delta$  receptors, and we conclude that these eight GABA<sub>A</sub>-Rs should be placed in category A as conclusively identified.

2. *Evidence from Other Ligands (GABA, Picrotoxin, General Anesthetics, and Ethanol) Confirms and Extends the List of Receptors.* Heterogeneity of GABA<sub>A</sub>-Rs was evident even before the cloning of the individual subunits as indicated by brain regional variation in radioligand binding, inhibition, and allosteric modulation involving the GABA, BZ, convulsant, and anesthetic sites as well as by photolabeling of GABA<sub>A</sub>-R subunits using radiolabeled

benzodiazepines (e.g., Sieghart, 1989; Olsen et al., 1990; Sieghart, 1995). Such evidence for subtypes was consistent with differential pharmacology assessed by electrophysiological recordings conducted using different cells (Macdonald and Olsen, 1994; Hevers and Lüddens, 1998). The potencies of GABA and antagonists such as gabazine (SR95531) vary slightly with the  $\alpha$  subunit (Bai et al., 2001), being more potent at  $\alpha 4$ , 5, and 6, than at  $\alpha 1$ , 2, and 3 subunit-containing receptors and with the  $\alpha 4\beta\delta$  and  $\alpha 6\beta\delta$  receptors showing even higher potency for GABA than the corresponding  $\gamma 2$  subunit-containing GABA<sub>A</sub>-R (Wallner et al., 2003). The  $\alpha 5\beta\gamma 2$ ,  $\alpha 4\beta\delta$ , and  $\alpha 6\beta\delta$  subtypes have recently been shown to be localized in extrasynaptic membranes, where they mediate a tonic inhibitory current, now recognized to have significant physiological relevance in controlling neuronal excitability, relevant to local circuitry and network activity (Bai et al., 2001; Caraiscos et al., 2004; Mody and Pearce, 2004; Farrant and Nusser, 2005). Many of these extrasynaptic GABA<sub>A</sub>-R show a high affinity for GABA and are resistant to desensitization, appropriate for their role in tonic inhibition. In addition, GABA has a low efficacy for these receptors, making them susceptible to higher efficacy for modulation by allosteric drugs such as nonbenzodiazepines and especially general anesthetics (Brown et al., 2002; Wohlfarth et al., 2002; Stell et al., 2003; Wallner et al., 2003; Caraiscos et al., 2004; Belelli et al., 2005). Overall, these drugs produce a larger charge transfer via extrasynaptic than synaptic receptors, and, in contrast to earlier dogma, the actions of some important drugs seem to involve modulation of extrasynaptic rather than synaptic receptors. Extrasynaptic GABA<sub>A</sub>-Rs show great heterogeneity with respect to subunit composition and include a significant fraction of the  $\gamma 2$ -containing GABA<sub>A</sub>-Rs, whereas those lacking  $\gamma 2$  are exclusively extrasynaptic.

The synaptic  $\gamma 2$  subunit-containing GABA<sub>A</sub>-Rs are all modulated by general anesthetics of diverse chemical structure including pentobarbital, etomidate, and propofol, as well as the neuroactive steroids, but the extrasynaptic  $\delta$  subunit-containing GABA<sub>A</sub>-Rs, and possibly the  $\alpha 5\beta\gamma 2$ , are more sensitive to these drugs (Brown et al., 2002; Wohlfarth et al., 2002; Stell et al., 2003; Wallner et al., 2003; Hemmings et al., 2005; Herd et al., 2007). As mentioned, the extrasynaptic  $\delta$  subunit-containing  $\alpha 4\beta\delta$  and  $\alpha 6\beta\delta$  GABA<sub>A</sub>-Rs show a significant modulation by nonbenzodiazepine drugs. Recordings in native tissues suggest therefore that both  $\gamma 2$ - and  $\delta$ -containing GABA<sub>A</sub>-Rs are clearly present and physiologically relevant. Glykys et al. (2007) demonstrated that the  $\delta$  subunit can pair with  $\alpha 1\beta$  in hippocampal interneurons, based on colocalization with immunostaining, electropharmacological properties, and changes in knockout mice. Although this single report is convincing, we decided that this subtype did not meet sufficient criteria to be included in the list of identified subtypes. Rather than omit it, the  $\alpha 1\beta\delta$  subtype is placed within a second category of native receptors

(category B, existence with high probability), so that readers are informed of receptors being considered for inclusion and can evaluate for themselves if they believe one qualifies.

The  $\gamma 2$  and  $\delta$  subunits may have selective  $\beta$  partners, but considerable evidence from colocalization, coimmunoprecipitation, and neuronal electrophysiology studies suggests that each partners with more than one  $\beta$  subunit (one at a time), so this provides a framework for additional subtypes. It was shown earlier that the nature of the  $\alpha$  subunit might affect modulator dose-response curves, but no effect was found for varying the  $\beta$  subunit (e.g., Hadingham et al., 1993; Thompson et al., 1996; Smith et al., 2004). Nevertheless, mutation of residues in the channel M2 domain of *any subunit*, but definitely showing some subunit selectivity in the details, can change modulator affinity or efficacy (Tierney et al., 1996; Hill-Venning et al., 1997; Thompson et al., 1999). Thus, mutations of residues in the M2 domain of  $\beta$  subunits can produce  $\beta$  subunit selectivity for pharmacological agents (Cestari et al., 1996; Rudolph and Antkowiak, 2004).

In addition, recent evidence suggests that the  $\beta$  subunit can affect both the pharmacological and channel properties of the GABA<sub>A</sub>-R, especially in the case of certain nonbenzodiazepine modulators and particularly for the extrasynaptic  $\delta$  subunit-containing receptors. Thus, several modulatory ligands, such as the nonbenzodiazepine loreclezole and chemically related ligands, such as the general anesthetic etomidate, show some selectivity for  $\beta 2$  and  $\beta 3$  over  $\beta 1$  subunit-containing receptors (Wingrove et al., 1994; Hill-Venning et al., 1997): these authors identified a single amino acid residue 15' in the M2 domain (counting from the intracellular N-terminal end of the channel-forming second transmembrane helix) responsible for this  $\beta$  subunit selectivity. Mutations of this residue were shown to affect sensitivity to the actions of etomidate in vitro (Belelli et al., 1997) and in vivo (Reynolds et al., 2003). A point mutation of the  $\beta 2$  subunit N265S that eliminates etomidate sensitivity in vitro loses the sedative-hypnotic effects of etomidate in the mouse knockin (Reynolds et al., 2003), whereas the  $\beta 3$  knockin N265M eliminates the immobilizing effect to a noxious stimulus of etomidate and suppresses loss of the righting reflex to this agent, typical of general anesthetics (Jurd et al., 2003).

The knockin mice showing either  $\beta 3$  or  $\beta 2$  insensitivity to etomidate in vivo allow detective work on which  $\beta$  subunit mediates the action of this drug on a given cell and current, and, along with other information about the physiology and subunit expression pattern in that cell, a good guess at the subunit composition of the functional receptors (e.g., which  $\alpha$  subunits go with which  $\beta$  subunit) (Rudolph and Antkowiak, 2004; Belelli et al., 2005; Herd et al., 2008). So far, however, the results did not provide evidence on whether the  $\gamma 2$ - or  $\delta$ -containing receptors, or both, mediate the etomidate

effects identified, although the  $\delta$ -containing receptors are more sensitive to modulation.

This residue 15' in the M2 "channel domain" of  $\beta$  subunits that affects etomidate sensitivity also affects sensitivity and to some degree  $\beta$  subunit selectivity of GABA<sub>A</sub>-R to other anesthetic modulators such as propofol, barbiturates, and volatile agents (Cestari et al., 1996; Rudolph and Antkowiak, 2004), consistent with the earlier observation that this residue in (presumably all)  $\alpha$  and  $\beta$  subunits affects sensitivity to volatile anesthetics and long-chain alcohols and may be part of a binding pocket for these drugs (Mihic et al., 1997; Ernst et al., 2005).

A few other ligands reported to show  $\beta$  selectivity include valerianic acid (Khom et al., 2007), some novel plant substances of a polyacetylene structure (Baur et al., 2005), tracazolol, and mefenamic acid (Korpi et al., 2002; Smith et al., 2004). This may be extended to ethanol (Wallner et al., 2003). The  $\delta$  subunit-containing GABA<sub>A</sub>-Rs, partnered with the  $\alpha 4$  or  $\alpha 6$  subunits and especially with  $\beta 3$ , are more sensitive than  $\gamma 2$ -containing receptors to general anesthetics, neurosteroids, GABA analogs such as gaboxadol (Brown et al., 2002; Wohlfarth et al., 2002; Chandra et al., 2006), and taurine (Hadley and Amin, 2007; Jia et al., 2008) as well as ethanol (Wallner et al., 2003; Hancher et al., 2005). The  $\alpha 4\beta 2\delta$  and  $\alpha 4\beta 3\delta$  receptors differ in sensitivity to modulators when recombinantly expressed in cells, and both clearly occur naturally, because some brain areas express  $\alpha 4$  and  $\delta$  but only  $\beta 2$  or  $\beta 3$  subunits and display a pharmacology that distinguishes between these receptors. Thus, thalamic relay nuclei mainly express the  $\beta 2$  subunit and the moderately ethanol-sensitive  $\alpha 4\beta 2\delta$  receptor definitely mediates the tonic current (Chandra et al., 2006). Dentate granule cells express high levels of highly ethanol-sensitive, presumably  $\alpha 4\beta 3\delta$  isoforms (Liang et al., 2006) but additionally express the etomidate-sensitive  $\alpha 4\beta 2\delta$  isoforms (Herd et al., 2008). This modulator selectivity for different  $\beta$  subunits in vitro and in vivo is supported by knockin mouse data. Partnering of either  $\beta 2$  or  $\beta 3$  with both the  $\alpha 4\delta$  and  $\alpha 6\delta$  subunits thus seems highly likely. To account for this additional evidence indicating the existence of  $\alpha 4\delta$  and  $\alpha 6\delta$  subunits combined with specific  $\beta$  subunits, we add  $\alpha 4\beta 2\delta$  and  $\alpha 4\beta 3\delta$ ,  $\alpha 6\beta 2\delta$  and  $\alpha 6\beta 3\delta$  subtypes in category A, replacing the generic  $\alpha 4\beta\delta$  and  $\alpha 6\beta\delta$  and bringing our total to 10. Subtypes dividing the  $\alpha\gamma 2$  subtypes depending on the  $\beta 2$  versus  $\beta 3$  partner, except for  $\alpha 1\beta 2\gamma 2$ , will not be listed until some compelling evidence is presented that the variants exist and differ in some property.

One strong candidate that nearly qualifies for category A is the  $\alpha 5\beta 3\gamma 2$  subtype. The  $\alpha 5$  and  $\beta 3$  subunits seem to be codepleted in mice lacking either the  $\beta 3$  or  $\alpha 5$  subunit (Olsen and Homanics, 2000), and the properties of recombinant  $\alpha 5\beta 3\gamma 2$  receptors seem to reflect those of a native subtype found in CA1 pyramidal neurons, cells enriched in both  $\alpha 5$  and  $\beta 3$  subunits, and distinct from

receptors in dentate gyrus granule cells (Burgard et al., 1996; Sur et al., 1998; McClennan and Twyman, 1999; Stell et al., 2003; Caraiscos et al., 2004). However, the specific partnering of  $\alpha 5$  with  $\beta 3$  has not been shown by coimmunoprecipitation or pharmacological changes in mutant mice, so we place this subtype in category B. Another example in this category is the  $\alpha 1\beta 3\gamma 2$  subtype. If we specify that  $\beta 2$  is the most common partner of  $\alpha 1\gamma 2$  and list it in category A, we need to ask whether  $\alpha 1$  can partner with other  $\beta$  subunits. Although evidence for  $\beta 1$  is lacking, the properties of  $\alpha 1\beta 3\gamma 2$  receptors in recombinant studies resemble those in certain neurons that express these subunits, some of which lack the  $\beta 2$  subunit (Whiting et al., 2000; Sieghart and Sperk, 2002). By contrast, the  $\beta$  partner with  $\alpha 2\gamma 2$  receptors is unknown; again the  $\alpha 2\beta 3\gamma 2$  is a likely possibility.

Although the  $\beta 1$  subunit is less abundant than the other  $\beta$  subunits, it is likely that some  $\beta 1$ -containing GABA<sub>A</sub>-Rs exist, based on regional distribution and coimmunoprecipitation data (Li and De Blas, 1997; Jechlinger et al., 1998). The preponderance of the evidence suggests that  $\beta 1$ -containing receptors exist. A potential pharmacological fingerprint, salicylidene salicylhydrazide, has been reported to be a negative allosteric modulator selective for  $\beta 1$  versus  $\beta 2$  or  $\beta 3$  subunit-containing receptors (Thompson et al., 2004). There is some evidence (Whiting et al., 2000) that cultured astrocytes express GABA<sub>A</sub>-Rs with  $\alpha 2\beta 1\gamma 1$  combinations, and they are uniquely enhanced by the BZ site ligand methy-6,7-dimethoxy-4-ethyl- $\beta$ -carboline-3-carboxylate (DMCM), a  $\beta$ -carboline. More evidence is needed to establish whether this subtype actually exists in brain. The  $\beta 1$  subunit probably partners with both  $\gamma$  or  $\delta$  subunits, but there is no conclusive evidence for either. At this time we will add just one subtype for  $\alpha\beta 1\gamma/\alpha\beta 1\delta$  in category B.

### C. Other, Nonligand, Delineation of Heterogeneity

#### 1. Subunit Composition.

*a. Minor subunits ( $\rho 1-3$ ,  $\gamma 1$ ,  $\gamma 3$ ,  $\varepsilon$ ,  $\theta$ , and  $\pi$ ).* First, evidence from all criteria supports the existence of at least one  $\rho$ -containing native receptor so we added  $\rho$  to the list in category A (Table 3), for a total of 11 identified receptors. That total is all that we find to meet sufficient criteria for this designation in 2008. However, each of the  $\rho$  subunit genes  $\rho 1$ ,  $\rho 2$ , and  $\rho 3$  probably encode at least one unique pharmacological subtype because their products have a differential brain distribution (Enz and Cutting, 1999; Greka et al., 2000). On the other hand, evidence for their separate formation of GABA-activated ion channels is scarce, because of the lack of antibodies for selective immunoprecipitation and the paucity of pharmacological tools suitable for their unequivocal identification as unique receptors. Hence, we place these three in a third category: C, tentative. However, we feel that the possible heteromerization of  $\rho$ , plus the possible combination with other GABA<sub>A</sub>-R or glycine receptor subunits (Pan et al., 2000; Hartmann et al., 2004; Mil-

ligan et al., 2004), does not qualify as subtypes for the list at this time.

The evidence for the existence of a native subtype containing any of the five minor subunits ( $\gamma 1$ ,  $\gamma 3$ ,  $\varepsilon$ ,  $\theta$ , and  $\pi$ ) is not as compelling as that for the other, more abundant, subunits ( $\alpha 1-6$ ,  $\beta 1-3$ ,  $\gamma 2$ , and  $\delta$ ) already described. This is attributable to a scarcity of studies and the lack of pharmacological tools and in vivo investigations. However, these subunits belong to the GABA<sub>A</sub>-R family and can form recombinant GABA-activated chloride channels on coexpression with other subunits, (Sieghart, 1995; Hevers and Lüddens, 1998) and exhibit differential regional distribution in the brain (Laurie et al., 1992; Persohn et al., 1992; Wisden et al., 1992; Davies et al., 1997; Hedblom and Kirkness, 1997; Whiting et al., 1997; Bonnert et al., 1999; Pirker et al., 2000; Moragues et al., 2000, 2002, 2003; Sinkkonen et al., 2000). Thus, evidence is tending toward all of them forming receptors in nature. In few cases are the possible heteropentameric subunit combinations known for these uncommon subunits (e.g., from immunoprecipitation studies using antibodies directed against  $\gamma 1$  or  $\gamma 3$  subunits) (Mossier et al., 1994; Quirk et al., 1994; Tögel et al., 1994) or  $\theta$  subunits (Bonnert et al., 1999) and thus the total number of subtypes may be large or small. Each subunit  $\gamma 1$ ,  $\gamma 3$ ,  $\varepsilon$ ,  $\theta$ , and  $\pi$  makes an unknown number of combinations with  $\alpha$  and  $\beta$  subunits, but probably will be slightly more than the minimum. In fact, the mere existence of these subunits has not been extended to show that the cells where they are expressed have any unique pharmacological or physiological properties produced from these subunits, although few serious searches have been attempted. Benzodiazepine insensitivity in the rat nucleus tractus solitarii correlates with  $\varepsilon$  expression (Kasparov et al., 2001), and prolongation of inhibitory postsynaptic currents recorded from cultured hypothalamic neurons has been shown to be negatively correlated with the detection of the  $\varepsilon$  subunit by RT-PCR (Sergeeva et al., 2005). In vitro recombinant studies suggest it should be possible to identify these rare subtypes if they exist. Because this has not been done yet, we decided to put these subtypes, combined with  $\alpha$  and  $\beta$ , in the third category, C, tentative. Unusual subunit combinations of these minor subunits remain possible. The  $\varepsilon$  subunit has been reported to partner with other combinations in recombinant expression (Davies et al., 1997, 2001; Moragues et al., 2000; Wagner et al., 2005; Jones and Henderson, 2007) but whether this occurs in nature has not been sufficiently investigated (Bollan et al., 2008). As mentioned above, "pathological" (non-native) subunit compositions can be forced to occur in recombinant expression systems. The partnership association of the  $\theta$  subunit has not been extensively studied (Bonnert et al., 1999; Sinkkonen et al., 2000) and,  $\pi$ , as a peripheral tissue constituent, has not been well characterized, although it forms channels in vitro with other GABA<sub>A</sub>-R subunits (Hedblom and Kirk-

TABLE 3  
GABA<sub>A</sub> receptor native oligomer list

Category A: Identified
$\alpha 1\beta 2\gamma 2$
$\alpha 2\beta\gamma 2$
$\alpha 3\beta\gamma 2$
$\alpha 4\beta\gamma 2$
$\alpha 4\beta 2\delta$
$\alpha 4\beta 3\delta$
$\alpha 5\beta\gamma 2$
$\alpha 6\beta\gamma 2$
$\alpha 6\beta 2\delta$
$\alpha 6\beta 3\delta$
$\rho$
Category B: Existence with high probability
$\alpha 1\beta 3\gamma 2$
$\alpha 1\beta\delta$
$\alpha 5\beta 3\gamma 2$
$\alpha\beta 1\gamma/\alpha\beta 1\delta$
$\alpha\beta$
$\alpha 1\alpha 6\beta\gamma/\alpha 1\alpha 6\beta\delta$
Category C: Tentative
$\rho 1$
$\rho 2$
$\rho 3$
$\alpha\beta\gamma 1$
$\alpha\beta\gamma 3$
$\alpha\beta\theta$
$\alpha\beta\epsilon$
$\alpha\beta\pi$
$\alpha\alpha\gamma\beta\gamma 2$

ness, 1997; Neelands and Macdonald, 1999; Korpi and Sinkkonen, 2006). Some of these subunits can be shown to express more than one type of channel in recombinant expression work. In these cases, it is particularly important to evaluate recombinant studies with caution unless the nature of the subunit composition of the expressed receptors is unambiguously determined.

*b.  $\alpha/\beta$  Pentamers.* Evidence has accumulated for the existence of receptors composed of  $\alpha$  and  $\beta$  subunits, only. First, Bencsits et al. (1999) demonstrated that a large proportion of the  $\alpha 4$  receptors (approximately 50%) are not associated with  $\gamma$  or  $\delta$  subunits and are possibly composed of  $\alpha$  and  $\beta$  subunits only. This is in contrast with  $\alpha 1$  receptors. Second, in several knockout mice it was demonstrated that receptors composed of only  $\alpha$  and  $\beta$  subunits do exist, for example, in  $\gamma 2$  knockout mice (Günther et al., 1995),  $\delta$  knockout mice (Tretter et al., 2001), or  $\alpha 1$  knockout mice (Ogris et al., 2006). This suggests that such receptors might develop under other pathological conditions also. It is possible that such receptors arise during epilepsy, hormone treatment, ethanol intake, or during development. Localization studies show that in several brain regions  $\alpha$  and  $\beta$  occur in the absence of  $\gamma$  and  $\delta$  subunits. Either these subunits have other functions or combine with other subunits or they form receptors of  $\alpha$  and  $\beta$  subunits only. Mortensen and Smart (2006) demonstrated by electrophysiological studies that there are extrasynaptic  $\alpha\beta$  receptors on rat hippocampal pyramidal neurons. We could not find evi-

dence for any specific  $\alpha$  or any specific  $\beta$  subunit in these  $\alpha\beta$  combinations or for the existence of more than one such subtype, so have placed one  $\alpha\beta$  subtype in category B (Table 3).

*c. Multiple  $\alpha$  or  $\beta$  (other?) isoforms per pentamer.* There is ample evidence from several different groups (Duggan et al., 1991; Lüddens et al., 1991; Khan et al., 1996; Sieghart and Sperk, 2002; Benke et al., 2004; reviewed in Whiting et al., 2000) for the coimmunoprecipitation of different  $\alpha$  or  $\beta$  subunits, whereas different  $\gamma$  subunits in most studies were observed not to coimmunoprecipitate to a significant extent (for discussion, see Sieghart and Sperk, 2002). These findings suggest there is no unspecific aggregation of different receptor subtypes, and a variety of controls were performed to demonstrate specific coprecipitation (Jechlinger et al., 1998; Pörtl et al., 2003). Nusser et al. (1998) demonstrated separate locations but also colocalization of  $\alpha 1$  and  $\alpha 6$  in the cerebellum, suggesting the possibility that the two  $\alpha$  subunits occur together in the same heteropentameric receptor. Benke et al. (2004) demonstrated the presence and abundance of GABA<sub>A</sub>-Rs containing two different types of  $\alpha$  subunits using point-mutated  $\alpha$  subunits and reported on the abundance of these receptors in an in vivo system: The  $\alpha 1$  predominantly forms  $\alpha 1\text{-}\alpha 1$ , whereas other subunits predominantly form hetero- $\alpha$ . Finally, Minier and Sigel (2004) described the differential properties of receptors containing  $\alpha 1$  and  $\alpha 6$  subunits in a defined arrangement in a recombinant system. Each receptor configuration has its own pharmacological signature. We decided that the  $\alpha 1\alpha 6\beta\gamma/\alpha 1\alpha 6\beta\delta$  qualified as a subtype for the list in category B. Furthermore, we placed an additional species of unknown subunit composition  $\alpha\alpha\gamma\beta\gamma 2$  in category C.

In addition, mixtures of  $\beta$  subunits probably do exist (Li and De Blas, 1997; Jechlinger et al., 1998), and two articles (Khan et al., 1994; Quirk et al., 1994) suggested that two kinds of  $\gamma$  subunit could be included. However, other researchers could not confirm the latter finding (Jechlinger et al., 1998; Pörtl et al., 2003).

*d. Receptors containing 1, 2, 4, or 5 different subunits.* The evidence for other combinations of subunits with other than three kinds of subunit (homo-oligomeric  $\beta 3$  subunit channels and hetero-oligomeric channels containing 2, 4, or 5 different subunits), except  $\rho$  and  $\alpha\beta$  as discussed, is limited to recombinant expression work and to some reports indicating the existence of two different  $\alpha$  or two different  $\beta$  subunits in the same native GABA<sub>A</sub>-R. None currently qualify for inclusion within the list (see discussion in Sieghart and Sperk, 2002).

*2. Localization, Trafficking, Post-Translational Modifications, and Associated Proteins.* Although these factors could alter receptor pharmacology (Kittler and Moss, 2003), little is known about such factors, no in vivo function has been identified unequivocally, and thus, none are accepted at this time.

## D. Tentative List of Naturally Occurring Receptor Subtypes

Three categories of entries are listed: identified, existence with high probability, and tentative.

## IV. Concluding Remarks

This classification starts with the 19 human GABA<sub>A</sub>-R genes (Table 1) and some evidence for the likely subunit combination and stoichiometry of the abundant ( $\alpha 1$ )<sub>2</sub>( $\beta 2$ )<sub>2</sub> $\gamma 2$  heteropentameric subtypes. Because of the emphasis of IUPHAR on the human clinic, the present discussion is limited to the human, rat, and mouse, acknowledging that GABA<sub>A</sub>-Rs exist in all organisms with a nervous system, including invertebrates. We further made the decision to concentrate on the usual location of GABA<sub>A</sub>-Rs, the central nervous system, realizing that there is expression of functional receptors in peripheral tissues.

In attempting to define native subtypes, we propose (Table 2) five criteria (two for recombinant studies and three for native studies) for inclusion of a given subunit combination on the list. We note that even for the 19 subunit genes, the evidence for some subunits participating in native subtypes varies considerably. Whereas there is some evidence that each of the 19 exists in at least one subtype, some of them meet few of the five criteria. We decided to use three different categories in making the list, consistent with the work in progress nature of such a list (Table 3). This is likewise the situation for the other members of the LGIC receptor superfamilies. The categories in the list are identified, existence with high probability, and tentative. We added the second category, existence with high probability, for those candidates that meet some but not a sufficient number of the criteria summarized here, allowing the reader to decide whether the evidence is convincing. Because the existence of native receptors containing the minor subunits did not meet even the criteria for this category but are likely to exist in at least one subtype each, we added the third category, tentative. One can see that each entry in the tentative category might actually exist as multiple discrete subtypes, and thus the number on the list will expand in the future, possibly by a substantial number, as suggested by Barnard et al. (1998). Our final list (Table 3) includes 11 subtypes in category A. We have 6 entries in category B. We placed 9 in category C, and if they all stay, that makes 26 in 2008, a number that is certain to increase over time.

## REFERENCES

- Atack JR (2005) The benzodiazepine binding site of GABA<sub>A</sub> receptors as a target for the development of novel anxiolytics. *Expert Opin Investig Drugs* **14**:601–618.
- Atack JR, Wafford KA, Tye SJ, Cook SM, Sohal B, Pike A, Sur C, Melillo D, Bristow L, Bromidge F, et al. (2006) TPA [7-(1,1-dimethyl)-6-(2-ethyl-2H-1,2,4-triazol-3-ylmethoxy)-3-(2-fluorophenyl)-1,2,4-triazolo[4,3-b]pyridazine], an agonist selective for  $\alpha 2$ - and  $\alpha 3$ -containing GABA<sub>A</sub> receptors, is a non-sedating anxiolytic in rodents and primates. *J Pharmacol Exp Ther* **316**:410–422.
- Bai D, Zhu G, Pennefather P, Jackson MF, MacDonald JF, and Orser BA (2001) Distinct functional and pharmacological properties of tonic and quantal inhibitory postsynaptic currents mediated by GABA<sub>A</sub> receptors in hippocampal neurons. *Mol Pharmacol* **59**:814–824.
- Barnard EA, Skolnick P, Olsen RW, Mohler H, Sieghart W, Biggio G, Braestrup C, Bateson AN, and Langer SZ (1998) International Union of Pharmacology. XV. Subtypes of  $\gamma$ -aminobutyric acid<sub>A</sub> receptors: classification on the basis of subunit structure and receptor function. *Pharmacol Rev* **50**:291–313.
- Barrera NP, Betts J, You H, Henderson RM, Martin IL, Dunn SMJ, and Edwardson JM (2008) Atomic force microscopy reveals the stoichiometry and subunit arrangement of the  $\alpha 4\beta 3\delta$  GABA<sub>A</sub> receptor. *Mol Pharmacol* **73**:960–967.
- Baumann SW, Baur R, and Sigel E (2002) Forced subunit assembly in  $\alpha 1\beta 2\gamma 2$  GABA<sub>A</sub> receptors: insight into the absolute arrangement. *J Biol Chem* **277**:46020–46025.
- Baur R, Simmen U, Senn M, Séquin U, and Sigel E (2005) Novel plant substances acting as  $\beta$  subunit isoform-selective positive allosteric modulators of GABA<sub>A</sub> receptors. *Mol Pharmacol* **68**:787–792.
- Belelli D, Lambert JJ, Peters JA, Wafford K, and Whiting PJ (1997) The interaction of the general anesthetic etomidate with the  $\gamma$ -aminobutyric acid type A receptor is influenced by a single amino acid. *Proc Natl Acad Sci U S A* **94**:11031–11036.
- Belelli D, Peden DR, Rosahl TW, Wafford KA, and Lambert JJ (2005) Extrasynaptic GABA<sub>A</sub> receptors of thalamocortical neurons: a molecular target for hypnotics. *J Neurosci* **25**:11513–11520.
- Ben-Ari Y (2002) Excitatory actions of GABA during development: the nature of the nurture. *Nat Rev Neurosci* **3**:728–739.
- Bencsits E, Ebert V, Tretter V, and Sieghart W (1999) A significant part of native  $\gamma$ -aminobutyric acid<sub>A</sub> receptors containing  $\alpha 4$  subunits do not contain  $\gamma$  or  $\delta$  subunits. *J Biol Chem* **274**:19613–19616.
- Benke D, Fakitsas P, Roggenmoser C, Michel C, Rudolph U, and Mohler H (2004) Analysis of the presence and abundance of GABA<sub>A</sub> receptors containing two different types of  $\alpha$  subunits in murine brain using point-mutated  $\alpha$  subunits. *J Biol Chem* **279**:43654–43660.
- Benke D, Mertens S, Trzeciak A, Gillessen D, and Mohler H (1991) GABA<sub>A</sub> receptors display association of  $\gamma 2$  subunits with  $\alpha 1$  and  $\beta 2/3$  subunits. *J Biol Chem* **266**:4478–4483.
- Bettler B, Kaufmann K, Mosbacher J, and Gassmann M (2004) Molecular structure and physiological functions of GABA<sub>B</sub> receptors. *Physiol Rev* **84**:835–867.
- Birnir B and Korpi ER (2007) The impact of sub-cellular location and intracellular neuronal proteins on properties of GABA<sub>A</sub> receptors. *Curr Pharm Des* **13**:3169–3177.
- Boehm SL 2nd, Ponomarev I, Jennings AW, Whiting PJ, Rosahl TW, Garrett EM, Blednov YA, and Harris RA (2004)  $\gamma$ -Aminobutyric acid A receptor subunit mutant mice: new perspectives on alcohol actions. *Biochem Pharmacol* **68**:1581–1602.
- Boileau AJ, Baur R, Sharkey LM, Sigel E, and Czajkowski C (2002) The relative amount of cRNA coding for  $\gamma 2$  subunits affects stimulation by benzodiazepines in GABA<sub>A</sub> receptors expressed in *Xenopus* oocytes. *Neuropharmacology* **43**:695–700.
- Boileau AJ, Pearce RA, and Czajkowski C (2005) Tandem subunits effectively constrain GABA<sub>A</sub> receptor stoichiometry and recapitulate receptor kinetics but are insensitive to GABA<sub>A</sub> receptor-associated protein. *J Neurosci* **25**:11219–11230.
- Bollan KA, Baur R, Hales TG, Sigel E, and Connolly CN (2008) The promiscuous role of the  $\epsilon$  subunit in GABA<sub>A</sub> receptor biogenesis. *Mol Cell Neurosci* **37**:610–621.
- Bonnert TP, McKernan RM, Farrar S, le Bourdellès B, Heavens RP, Smith DW, Hewson L, Rigby MR, Sirinathsinghji DJ, Brown N, Wafford KA, and Whiting PJ (1999)  $\theta$ , a novel  $\gamma$ -aminobutyric acid type A receptor subunit. *Proc Natl Acad Sci U S A* **96**:9891–9896.
- Borboni P, Porcio O, Fusco A, Sesti G, Lauro R, and Marlier LN (1994) Molecular and cellular characterization of the GABA<sub>A</sub> receptor in the rat pancreas. *Mol Cell Endocrinol* **103**:157–163.
- Bormann J and Feigenspan A (1995) GABA<sub>C</sub> receptors. *Trends Neurosci* **18**:515–519.
- Bowery NG, Bettler W, Froestl W, Gallager JP, Marshall F, Raiteri M, Bonner TI, and Enna SJ (2002) International Union Pharmacology. XXXIII. Mammalian  $\gamma$ -aminobutyric acid<sub>B</sub> receptors: structure and function. *Pharmacol Rev* **54**:247–264.
- Breitinger HG and Becker CM (2002) The inhibitory glycine receptor—simple views of a complicated channel. *ChemBiochem* **3**:1042–1052.
- Brejci K, van Dijk WJ, Klaassen RV, Schuurmans M, van Der Oost J, Smit AB, and Sixma TK (2001) Crystal structure of an ACh-binding protein reveals the ligand-binding domain of nicotinic receptors. *Nature* **411**:269–276.
- Brown N, Kerby J, Bonnert TP, Whiting PJ, and Wafford KA (2002) Pharmacological characterization of a novel cell line expressing human  $\alpha 4\beta 3\delta$  GABA<sub>A</sub> receptors. *Br J Pharmacol* **136**:965–974.
- Buckingham SD, Biggin PC, Sattelle BM, Brown LA, and Sattelle DB (2005) Insect GABA receptors: splicing, editing, and targeting by antiparasitics and insecticides. *Mol Pharmacol* **68**:942–951.
- Burgard EC, Tietz EI, Neelands TR, and Macdonald RL (1996) Properties of recombinant GABA<sub>A</sub> receptor isoforms containing the  $\alpha 5$  subunit subtype. *Mol Pharmacol* **50**:119–127.
- Caraiscos VB, Elliott EM, You-Ten KE, Cheng VY, Belleli D, Newell JG, Jackson MF, Lambert JJ, Rosahl TW, Wafford KA, et al. (2004) Tonic inhibition in mouse hippocampal CA1 pyramidal neurons is mediated by  $\alpha 5$  subunit containing GABA<sub>A</sub> receptors. *Proc Natl Acad Sci U S A* **101**:3662–3667.
- Cestari IN, Uchida I, Li L, Burt D, and Yang J (1996) The agonist action of pentobarbital on GABA<sub>A</sub>  $\beta$  subunit homomeric receptors. *Neuroreport* **7**:943–947.
- Chandra D, Jia F, Liang J, Peng Z, Suryanarayanan A, Werner DF, Spigelman I, Houser CR, Olsen RW, Harrison NL, et al. (2006) GABA<sub>A</sub> receptor  $\alpha 4$  subunits mediate extrasynaptic inhibition in thalamus and dentate gyrus and are required for the action of gaboxadol. *Proc Natl Acad Sci U S A* **103**:15230–15235.
- Chen PE and Wyllie DJ (2006) Pharmacological insights obtained from structure-function studies of ionotropic glutamate receptors. *Br J Pharmacol* **147**:839–853.
- Cheng VY, Martin LJ, Elliott EM, Kim JH, Mount HT, Taverna FA, Roder JC, MacDonald JF, Bhabri A, Collinson N, et al. (2006)  $\alpha 5$ GABA<sub>A</sub> receptors mediate the amnesic but not sedative-hypnotic effects of the general anesthetic etomidate. *J Neurosci* **26**:3713–3720.
- Collinson N, Cothliff R, Rosahl TW, Sur C, Kuenzi F, Howell O, Seabrook GR, Atack

- JR, McKernan RM, Dawson GR, et al. (2000) Role of the  $\alpha 5$  subunit of the GABA<sub>A</sub> receptor in learning and memory. *Eur J Neurosci* **12** (Suppl 11):171.
- Corringer PJ, Le Novère N, and Changeux JP (2000) Nicotinic receptors at the amino acid level. *Annu Rev Pharmacol Toxicol* **40**:431–458.
- Crestani F, Keist R, Fritschy JM, Benke D, Vogt K, Prut L, Blüthmann H, Möhler H, and Rudolph U (2002) Trace fear conditioning involves hippocampal  $\alpha 5$  GABA<sub>A</sub> receptors. *Proc Natl Acad Sci U S A* **99**:8980–8985.
- Crestani F, Martin JR, Möhler H, and Rudolph U (2000) Resolving differences in GABA<sub>A</sub> receptor mutant mouse studies. *Nat Neurosci* **3**:1059.
- Criswell HE, McCown TJ, Moy SS, and Breese G (1997) Action of zolpidem on responses to GABA in relation to mRNAs for GABA<sub>A</sub> receptor  $\alpha$  subunits within single cells: evidence for multiple functional GABA<sub>A</sub> isoreceptors on individual neurons. *Neuropharmacology* **36**:1641–1652.
- Cromer BA, Morton CJ, and Parker MW (2002) Anxiety over GABA<sub>A</sub> receptor structure relieved by AChBP. *Trends Biochem Sci* **27**:280–287.
- Davies PA, Hanna MC, Hales TG, and Kirkness EF (1997) Insensitivity to anaesthetic agents conferred by a class of GABA<sub>A</sub> receptor subunit. *Nature* **385**:820–823.
- Davies PA, Kirkness EF, and Hales TG (2001) Evidence for the formation of functionally distinct  $\alpha\beta\gamma\epsilon$  GABA<sub>A</sub> receptors. *J Physiol* **537**:101–113.
- Davies PA, Pistis M, Hanna MC, Peters JA, Lambert JJ, Hales TG, and Kirkness EF (1999) The 5-HT<sub>3B</sub> subunit is a major determinant of serotonin receptor function. *Nature* **397**:359–363.
- Davies PA, Wang W, Hales TG, and Kirkness EF (2003) A novel class of ligand-gated ion channel is activated by Zn<sup>2+</sup>. *J Biol Chem* **278**:712–717.
- Dawson GR, Maubach KA, Collinson N, Cobain M, Everitt BJ, MacLeod AM, Choudhury HI, McDonald LM, Pillai G, Rycroft W, et al. (2006) An inverse agonist selective for  $\alpha 5$  subunit-containing GABA<sub>A</sub> receptors enhances cognition. *J Pharmacol Exp Ther* **316**:1335–1345.
- Dellisanti CH, Yao Y, Stroud JC, Wang ZZ, and Chen L (2007) Crystal structure of the extracellular domain of nAChR  $\alpha 1$  bound to  $\alpha$ -bungarotoxin at 1.94 Å resolution. *Nat Neurosci* **10**:953–962.
- Dias R, Sheppard WF, Fradley RL, Garrett EM, Stanley JL, Tye SJ, Goodacre S, Lincoln RJ, Cook SM, Conley R, et al. (2005) Evidence for a significant role of  $\alpha 3$ -containing GABA<sub>A</sub> receptors in mediating the anxiolytic effects of benzodiazepines. *J Neurosci* **25**:10682–10688.
- Duggan MJ, Pollard S, and Stephenson FA (1991) Immunoaffinity purification of GABA<sub>A</sub> receptor  $\alpha$ -subunit iso-oligomers: demonstration of receptor populations containing  $\alpha 1\alpha 2$ ,  $\alpha 1\alpha 3$ , and  $\alpha 2\alpha 3$  subunit pairs. *J Biol Chem* **266**:24778–24784.
- Enz R and Cutting GR (1999) GABA<sub>C</sub> receptor rho subunits are heterogeneously expressed in the human CNS and form homo- and hetero-oligomers with distinct physical properties. *Eur J Neurosci* **11**:41–50.
- Erickson SS and Boileau AJ (2007) Tandem cointure: Cys-loop receptor concatemer insights and caveats. *Mol Neurobiol* **35**:113–128.
- Ernst M, Brauchart D, Borech S, and Sieghart W (2003) Comparative modeling of GABA<sub>A</sub> receptors: limits, insights, future developments. *Neuroscience* **119**:933–943.
- Ernst M, Bruckner S, Borech S, and Sieghart W (2005) Comparative models of GABA<sub>A</sub> receptor extracellular and transmembrane domains: important insights into pharmacology and function. *Mol Pharmacol* **68**:1291–1300.
- Farrant M and Nusser Z (2005) Variations on an inhibitory theme: phasic and tonic activation of GABA<sub>A</sub> receptors. *Nat Rev Neurosci* **6**:215–229.
- Fradley RL, Guscott MR, Bull S, Hallett DJ, Goodacre SC, Wafford KA, Garrett EM, Newman RJ, O'Meara GF, Whiting PJ, et al. (2007) Differential contribution of GABA<sub>A</sub> receptor subtypes to the anticonvulsant efficacy of benzodiazepine site ligands. *J Psychopharmacol* **21**:384–391.
- Fritschy JM and Möhler H (1995) GABA<sub>A</sub> receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits. *J Comp Neurol* **359**:154–194.
- Galzi JL and Changeux JP (1994) Ligand-gated ion channels as unconventional allosteric proteins. *Curr Opin Struct Biol* **4**:554–565.
- Gardner CR, Tully WR, and Hedgecock CJ (1993) The rapidly expanding range of neuronal benzodiazepine receptor ligands. *Prog Neurobiol* **40**:1–61.
- Glykys J, Peng Z, Chandra D, Homanics GE, Houser CR, and Mody I (2007) A new naturally occurring GABA<sub>A</sub> receptor subunit partnership with high sensitivity to ethanol. *Nature Neurosci* **10**:40–48.
- Greka A, Lipton SA, and Zhang D (2000) Expression of GABA<sub>C</sub> receptor  $\rho 1$  and  $\rho 2$  subunits during development of the mouse retina. *Eur J Neurosci* **12**:3575–3582.
- Günther U, Benson J, Benke D, Fritschy JM, Reyes G, Knoflach F, Crestani F, Aguzzi A, Arigoni M, Lang Y, et al. (1995) Benzodiazepine-insensitive mice generated by targeted disruption of the  $\gamma 2$  subunit gene of  $\gamma$ -aminobutyric acid type A receptors. *Proc Natl Acad Sci U S A* **92**:7749–7753.
- Gutiérrez A, Khan ZU, and De Blas AL (1994) Immunocytochemical localization of  $\gamma 2$  short and  $\gamma 2$  long subunits of the GABA<sub>A</sub> receptor in the rat brain. *J Neurosci* **14**:7168–7179.
- Gutiérrez A, Khan ZU, Miralles CP, and De Blas AL (1996) Altered expression of  $\gamma 2L$  and  $\gamma 2S$  GABA<sub>A</sub> receptor subunits in the aging rat brain. *Brain Res Mol Brain Res* **35**:91–102.
- Hadingham KL, Wingrove PB, Wafford KA, Bain C, Kemp JA, Palmer KJ, Wilson AW, Wilcox AS, Sikela JM, and Ragan CI (1993) Role of the  $\beta$  subunit in determining the pharmacology of human  $\gamma$ -aminobutyric acid type A receptors. *Mol Pharmacol* **44**:1211–1218.
- Hadley SH and Amin J (2007) Rat  $\alpha 6\beta 2\delta$  GABA<sub>A</sub> receptors exhibit two distinct and separable agonist affinities. *J Physiol* **581**:1001–1018.
- Hanchar HJ, Chutrinopkun P, Meera P, Sieghart W, Supavilai P, Wallner M, and Olsen RW (2006) Ethanol potently and competitively inhibits the binding of the alcohol antagonist Ro15-4513 to  $\alpha 4\beta 3\delta$  GABA<sub>A</sub> receptors. *Proc Natl Acad Sci U S A*, **103**:8546–8551.
- Hanchar HJ, Dodson PD, Olsen RW, Otis TS, and Wallner M (2005) Alcohol induced motor impairment caused by increased extrasynaptic GABA<sub>A</sub> receptor activity. *Nat Neurosci* **8**:339–345.
- Hartmann K, Stief F, Draguhn A, and Frahm C (2004) Ionotropic GABA receptors with mixed pharmacological properties of GABA<sub>A</sub> and GABA<sub>C</sub> receptors. *Eur J Pharmacol* **497**:139–146.
- Hedblom E and Kirkness EF (1997) A novel class of GABA<sub>A</sub> receptor subunit in tissues of the reproductive system. *J Biol Chem* **272**:15346–15350.
- Hemmings HC, Akabas MH, Goldstein PA, Trudell JR, Orser BA, and Harrison NL (2005) Emerging molecular mechanisms of general anesthetic action. *Trends Pharmacol Sci* **26**:503–510.
- Herd MB, Belelli D, and Lambert JJ (2007) Neurosteroid modulation of synaptic and extrasynaptic GABA<sub>A</sub> receptors. *Pharmacol Ther* **116**:20–34.
- Herd MB, Haythornthwaite AR, Rosahl TW, Wafford KW, Homanics GE, Lambert JJ, and Belelli D (2008) The expression of GABA<sub>A</sub>  $\beta$  subunit isoforms in synaptic and extrasynaptic receptor populations of mouse dentate gyrus granule cells. *J Physiol* **586**:989–1004.
- Hevers W and Lüddens H (1998) The diversity of GABA<sub>A</sub> receptors: pharmacological and electrophysiological properties of GABA<sub>A</sub> channel subtypes. *Mol Neurobiol* **18**:35–86.
- Hevers W, Korpi ER, and Lüddens H (2000) Assembly of functional  $\alpha 6\beta 3\gamma 2\delta$  GABA<sub>A</sub> receptors in vitro. *Neuroreport* **11**:4103–4106.
- Hill-Venning C, Belelli D, Peters JA, and Lambert JJ (1997) Subunit-dependent interactions of the general anaesthetic etomidate with the GABA<sub>A</sub> receptor. *Br J Pharmacol* **120**:749–756.
- Im WB, Pregenzer JF, Binder JA, Dillon GH, and Alberts GL (1995) Chloride channel expression with the tandem construct of  $\alpha 6$ - $\beta 2$  GABA<sub>A</sub> receptor subunit requires a monomeric subunit of  $\alpha 6$  or  $\gamma 2$ . *J Biol Chem* **270**:26063–26066.
- Itier V, Depoortere H, Scatton B, and Avenet P (1996) Zolpidem functionally discriminates subtypes of native GABA<sub>A</sub> receptors in acutely dissociated rat striatal and cerebellar neurons. *Neuropharmacology* **35**:137–145.
- Jechlinger M, Pelz R, Tretter V, Klausberger T, and Sieghart W (1998) Subunit composition and quantitative importance of hetero-oligomeric receptors: GABA<sub>A</sub> receptors containing  $\alpha 6$  subunits. *J Neurosci* **18**:2449–2457.
- Jia F, Yue M, Chandra D, Keramidias A, Goldstein PA, Homanics GE, and Harrison NL (2008) Taurine is a potent activator of extrasynaptic GABA<sub>A</sub> receptors in the thalamus. *J Neurosci* **28**:106–115.
- Johnston GAR (1996) GABA<sub>C</sub> receptors: relatively simple transmitter-gated ion channels? *Trends Pharmacol Sci* **17**:319–323.
- Johnston GAR (2002) Medicinal chemistry and molecular pharmacology of GABA<sub>C</sub> Receptors. *Curr Top Med Chem* **2**:903–913.
- Johnston GAR (2005) GABA<sub>A</sub> receptor channel pharmacology. *Curr Pharm Des* **11**:1867–1885.
- Jones A, Korpi ER, McKernan RM, Pelz R, Nusser Z, Makela R, Mellor JR, Pollard S, Bahn S, Stephenson FA, et al. (1997) Ligand-gated ion channel subunit partnerships: GABA<sub>A</sub> receptor  $\alpha 6$  subunit gene inactivation inhibits delta subunit expression. *J Neurosci* **17**:1350–1362.
- Jones BL and Henderson LP (2007) Trafficking and potential assembly patterns of  $\epsilon$ -containing GABA<sub>A</sub> receptors. *J Neurochem* **103**:1258–1271.
- Jurd R, Arras M, Lambert S, Drexler B, Sieghart W, Crestani F, Zaugg M, Vogt KE, Ledermann B, Antkowiak B, et al. (2003) General anesthetic actions in vivo strongly attenuated by a point mutation in the GABA<sub>A</sub> receptor beta3 subunit. *FASEB J* **17**:250–252.
- Kaila K, Lamsa K, Smirnov S, Taira T, and Voipio J (1997) Long-lasting GABA-mediated depolarization evoked by high-frequency stimulation in pyramidal neurons of rat hippocampal slice is attributable to a network-driven, bicarbonate-dependent K<sup>+</sup> transient. *J Neurosci* **17**:7662–7672.
- Kasparov S, Davies KA, Patel UA, Boscan P, Garret M, and Paton JF (2001) GABA<sub>A</sub> receptor epsilon subunit may confer benzodiazepine insensitivity to the caudal aspect of the nucleus tractus solitarius of the rat. *J Physiol* **536**:785–796.
- Keramidas A, Moorhouse AJ, Schofield PRP, and Barry PH (2004) Ligand-gated ion channels: mechanisms underlying ion selectivity. *Prog Biophys Mol Biol* **86**:161–204.
- Khakh BS, Burnstock G, Kennedy C, King BF, North RA, Seguela P, Voigt M, and Humphrey PP (2001) International Union of pharmacology. XXIV. Current status of the nomenclature and properties of P2X receptors and their subunits. *Pharmacol Rev* **53**:107–118.
- Khakh BS and North RA (2006) P2X receptors as cell-surface ATP sensor in health and disease. *Nature* **442**:3:527–532.
- Khan Z-U, Gutiérrez A, and De Blas AL (1994) Short and long form  $\gamma 2$  subunits of the GABA<sub>A</sub>/benzodiazepine receptors. *J Neurochem* **63**:1466–1476.
- Khan Z-U, Gutiérrez A, and De Blas AL (1996) The  $\alpha 1$  and  $\alpha 6$  subunits can coexist in the same cerebellar GABA<sub>A</sub> receptor maintaining their individual benzodiazepine binding specificities. *J Neurochem* **66**:685–691.
- Khom S, Baburin I, Timin EN, Hohaus A, Sieghart W, and Hering S (2006) Pharmacological properties of GABA<sub>A</sub> receptors containing  $\gamma 1$  subunits. *Mol Pharmacol* **69**:640–649.
- Khom S, Baburin I, Timin E, Hohaus A, Trauner G, Kopp B, and Hering S (2007) Valerianic acid potentiates and inhibits GABA<sub>A</sub> receptors: molecular mechanism and subunit specificity. *Neuropharmacology* **53**:178–187.
- Kittler JT and Moss SJ (2003) Modulation of GABA<sub>A</sub> receptor activity by phosphorylation and receptor trafficking: implications for the efficacy of synaptic inhibition. *Curr Opin Neurobiol* **13**:341–347.
- Klausberger T, Roberts JDB, and Somogyi P (2002) Cell type and input-specific differences in the number and subtypes of synaptic GABA<sub>A</sub> receptors in the hippocampus. *J Neurosci* **22**:2513–2521.
- Kofuji P, Wang JB, Moss SJ, Hagan RL, and Burt DR (1991) Generation of two forms of the gamma-aminobutyric acidA receptor  $\gamma 2$ -subunit in mice by alternative splicing. *J Neurochem* **56**:713–715.
- Korpi ER, Gründer G, and Lüddens H (2002) Drug interactions at GABA<sub>A</sub> receptors. *Prog Neurobiol* **67**:113–159.
- Korpi ER and Sinkkonen ST (2006) GABA<sub>A</sub> receptor subtypes as targets for neuropsychiatric drug development. *Pharmacol Ther* **109**:12–32.
- Laurie DJ, Wisden W, and Seeburg PH (1992) The distribution of 13 GABA<sub>A</sub> receptor



- subunit mRNAs in the rat brain. III. Embryonic and postnatal development. *J Neurosci* **12**:4151–4172.
- Lewis TM, Harkness PC, Sivilotti LG, Colquhoun D, and Millar NS (1997) The ion channel properties of a rat recombinant neuronal nicotinic receptor are dependent on the host cell type. *J Physiol* **505**:299–306.
- Li M and De Blas A (1997) Coexistence of two  $\beta$  subunit isoforms in the same  $\gamma$ -aminobutyric acid type A receptor. *J Biol Chem* **272**:16564–16569.
- Liang J, Suryanarayanan A, Chandra D, Homanics GE, Olsen RW, and Spigelman I (2008) Functional consequences of GABA<sub>A</sub> receptor  $\alpha 4$  subunit deletion on synaptic and extrasynaptic currents in mouse dentate granule cells. *Alcohol Clin Exp Res* **32**:19–26.
- Liang J, Zhang N, Cagetti E, Houser CR, Olsen RW, and Spigelman I (2006) Chronic intermittent ethanol-induced switch of ethanol actions from extrasynaptic to synaptic hippocampal GABA<sub>A</sub> receptors. *J Neurosci* **26**:1749–1758.
- Löw K, Crestani F, Keist R, Benke D, Brünig I, Benson JA, Fritschy JM, Rüllicke T, Blüthmann H, Möhler H, et al. (2000) Molecular and neuronal substrate for the selective attenuation of anxiety. *Science* **290**:131–134.
- Lüddens H, Killisch I, and Seeburg PH (1991) More than one  $\alpha$  variant may exist in a GABA<sub>A</sub>/benzodiazepine receptor complex. *J Recept Res* **11**:535–551.
- Lukas RJ and Bencherif M (2006) Recent developments in nicotinic acetylcholine receptor biology, in *Biological and Biophysical Aspects of Ligand-gated Ion Channel Receptor Superfamilies* (Arias HR eds) pp 2–33, Research Signpost, Kerala, India.
- Macdonald RL and Olsen RW (1994) GABA<sub>A</sub> receptor channels. *Annu Rev Neurosci* **17**:569–602.
- Martin DL and Olsen RW, editors (2000) *GABA in the Nervous System: The View at 50 Years*. Lippincott, Williams & Wilkins, New York.
- Marty A and Liano I (2005) Excitatory effects of GABA in established brain networks. *Trends Neurosci* **28**:284–289.
- McClelland AML and Twyman RE (1999) Receptor system response kinetics reveal functional subtypes of native murine and recombinant human GABA<sub>A</sub> receptors. *J Physiol* **515**:717–727.
- McKernan RM, Rosahl TW, Reynolds DS, Sur C, Wafford KA, Atack JR, Farrar S, Myers J, Cook G, Ferris P, Garrett L, Bristow L, Marshall G, Macaulay A, Brown N, Howell O, Moore KW, Carling RW, Street LJ, Castro JL, Ragan CI, Dawson GR, and Whiting PJ (2000) Sedative but not anxiolytic properties of benzodiazepines are mediated by the GABA<sub>A</sub> receptor  $\alpha 1$  subtype. *Nat Neurosci* **3**:587–592.
- Meier J and Grantyn R (2004) Preferential accumulation of GABA<sub>A</sub> receptor  $\gamma 2L$ , not  $\gamma 2S$ , cytoplasmic loops at rat spinal cord inhibitory synapses. *J Physiol* **559**:355–365.
- Mihalek RM, Bowers BJ, Wehner JM, Kralic JE, VanDoren MJ, Morrow AL, and Homanics GE (2001) GABA<sub>A</sub>-receptor  $\delta$  subunit knockout mice have multiple defects in behavioral responses to ethanol. *Alcohol Clin Exp Res* **25**:1708–1718.
- Mihic SJ, Ye Q, Wick MJ, Koltchine VV, Krasowski MD, Finn SE, Mascia MP, Valenzuela CF, Hanson KK, Greenblatt EP, Harris RA, and Harrison NL (1997) Sites of alcohol and volatile anesthetic action on GABA<sub>A</sub> and glycine receptors. *Nature* **389**:385–389.
- Milligan CJ, Buckley NJ, Garret M, Deuchars J, and Deuchars SA (2004) Evidence for inhibition mediated by co-assembly of GABA<sub>A</sub> and GABA<sub>C</sub> receptor subunits in native central neurons. *J Neurosci* **24**:7241–7250.
- Minier F and Sigel E (2004) Positioning of the  $\alpha$ -subunit isoforms confers a functional signature to gamma-aminobutyric acid type A receptors. *Proc Natl Acad Sci U S A* **10**:10–15.
- Miyazawa A, Fujiyoshi Y, and Unwin N (2003) Structure and gating mechanism of the acetylcholine receptor pore. *Nature* **423**:949–955.
- Mody I and Pearce RA (2004) Diversity of inhibitory neurotransmission through GABA<sub>A</sub> receptors. *Trends Neurosci* **27**:569–575.
- Möhler H (2007) Molecular recognition of cognitive functions and developmental plasticity: impact of GABA<sub>A</sub> receptors. *J Neurochem* **102**:1–12.
- Moragues N, Ciofi P, Lafon P, Odessa MF, Tramu G, and Garret M (2000) cDNA cloning and expression of a  $\gamma$ -aminobutyric acid A receptor  $\epsilon$ -subunit in rat brain. *Eur J Neurosci* **12**:4318–4330.
- Moragues N, Ciofi P, Lafon P, Tramu G, and Garret M (2003) GABA<sub>A</sub> receptor  $\epsilon$  subunit expression in identified peptidergic neurons of the rat hypothalamus. *Brain Res* **967**:285–289.
- Moragues N, Ciofi P, Tramu G, and Garret M (2002) Localisation of GABA<sub>A</sub> receptor  $\epsilon$ -subunit in cholinergic and aminergic neurons and evidence for co-distribution with the  $\theta$ -subunit in rat brain. *Neuroscience* **111**:657–669.
- Morris HV, Dawson GR, Reynolds DS, Atack JR, Stephens DN (2006) Both  $\alpha 2$  and  $\alpha 3$  GABA<sub>A</sub> receptor subtypes mediate the anxiolytic properties of benzodiazepine site ligands in the conditioned emotional response paradigm. *Eur J Neurosci* **23**:2495–2504.
- Mortensen M and Smart TG (2006) Extrasynaptic  $\alpha/\beta$  subunit GABA<sub>A</sub> receptors on rat hippocampal pyramidal neurons. *J Physiol* **577**:841–856.
- Moser N, Mechawar N, Jones I, Gochberg-Sarver A, Orr-Urtreger A, Plomann M, Saalas R, Molles B, Marubio L, Roth U, et al. (2007) Evaluating the suitability of nicotinic acetylcholine receptor antibodies for standard immunodetection procedures. *J Neurochem* **102**:479–492.
- Mossier B, Tögel M, Fuchs K, and Sieghart W (1994) Immunoaffinity purification of  $\gamma$ -aminobutyric acid<sub>A</sub> (GABA<sub>A</sub>) receptors containing gamma1-subunits: evidence for the presence of a single type of  $\gamma$ -subunit in GABA<sub>A</sub> receptors. *J Biol Chem* **269**:25777–25782.
- Neelands TR and Macdonald RL (1999) Incorporation of the  $\pi$  subunit into functional GABA<sub>A</sub> receptors. *Mol Pharmacol* **56**:598–610.
- Nusser Z, Sieghart W, and Somogyi P (1998) Segregation of different GABA<sub>A</sub> receptors to synaptic and extrasynaptic membranes of cerebellar granule cells. *J Neurosci* **18**:1693–1703.
- Ogris W, Lehner R, Fuchs K, Furtmüller B, Hoyer H, Homanics GE, and Sieghart W (2006) Investigation of the abundance and subunit composition of GABA<sub>A</sub> receptor subtypes in the cerebellum of  $\alpha 1$  subunit-deficient mice. *J Neurochem* **96**:136–147.
- Ohlson J, Pedersen JS, Haussler D, and Ohman M (2007) Editing modifies the GABA<sub>A</sub> receptor subunit  $\alpha 3$ . *RNA* **13**:698–703.
- Olsen RW and Gordey M (2000) GABA<sub>A</sub> receptor chloride ion channels, in *Handbook of Experimental Pharmacology*, vol 147, *Pharmacology of Ionic Channel Function: Activators and Inhibitors* (Endo M, Kurachi Y, Mishina M eds) pp 499–517, Springer Verlag, Heidelberg.
- Olsen RW, Hancher HJ, Meera P, and Wallner M (2007) GABA<sub>A</sub> receptor subtypes: the 'one glass of wine' receptors. *Alcohol* **41**:201–209.
- Olsen RW and Homanics GE (2000) Function of GABA<sub>A</sub> receptors: insights from mutant and knockout mice, in *GABA in the Nervous System: The View at 50 Years* (Martin DL and Olsen RW eds) pp. 81–96. Philadelphia, Lippincott, Williams & Wilkins.
- Olsen RW, McCabe RT, and Wamsley JK (1990) GABA<sub>A</sub> receptor subtypes: autoradiographic comparison of GABA, benzodiazepine and convulsant binding sites in the rat central nervous system. *J Chem Neuroanat* **3**:59–76.
- Olsen RW and Sawyer GW (2004) GABA<sub>A</sub> receptor, in *Encyclopedia of Biological Chemistry*, vol 2, pp 162–166, Elsevier, New York.
- Pan Y and Qian H (2005) Interactions between  $\rho$  and  $\gamma 2$  subunits of the GABA receptor. *J Neurochem* **94**:482–490.
- Pan ZH, Zhang D, Zhang X, and Lipton SA (2000) Evidence for coassembly of mutant GABA<sub>C</sub>  $\rho 1$  with GABA<sub>A</sub>  $\gamma 2S$ , glycine  $\alpha 1$  and glycine  $\alpha 2$  receptor subunits in vitro. *Eur J Neurosci* **12**:3137–3145.
- Peng Z, Hauer B, Mihalek RM, Homanics GE, Sieghart W, Olsen RW, and Houser CR (2002) GABA<sub>A</sub> receptor subunit changes in  $\delta$  subunit-deficient mice: altered expression of  $\alpha 4$  and  $\gamma 2$  subunits in the forebrain. *J Comp Neurol* **446**:179–197.
- Persohn E, Malherbe P, and Richards JG (1991) In situ hybridization histochemistry reveals a diversity of GABA<sub>A</sub> receptor subunit mRNAs in neurons of the rat spinal cord and dorsal root ganglia. *Neuroscience* **42**:497–507.
- Persohn E, Malherbe P, and Richards JG (1992) Comparative molecular neuroanatomy of cloned GABA<sub>A</sub> receptor subunits in the rat CNS. *J Comp Neurol* **326**:193–216.
- Peters JA, Hales TG, and Lambert JJ (2005) Molecular determinants of single-channel conductance and ion selectivity in the Cys-loop family: insights from the 5-HT<sub>3</sub> receptor. *Trends Pharmacol Sci* **26**:587–594.
- Pin JP, Neubig R, Bouvier M, Devi L, Filizola M, Javitch JA, Lohse MJ, Milligan G, Palczewski K, Parmentier M, et al. (2007) International Union of Basic and Clinical Pharmacology. LXVII. Recommendations for the recognition and nomenclature of G protein-coupled receptor heteromultimers. *Pharmacol Rev* **59**:5–13.
- Pirker S, Schwarzer C, Wieselthaler A, Sieghart W, and Sperk G (2000) GABA<sub>A</sub> receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience* **101**:815–833.
- Pörtl A, Hauer B, Fuchs K, Tretter V, and Sieghart W (2003) Subunit composition and quantitative importance of GABA<sub>A</sub> receptor subtypes in the cerebellum of mouse and rat. *J Neurochem* **87**:1444–1455.
- Ponomarev I, Maiya R, Harnett MT, Schafer GL, Ryabinin AE, Blednov Y, Morikawa H, Boehm SL 2nd, Homanics GE, Berman A, et al. (2006) Transcriptional signatures of cellular plasticity in mice lacking the  $\alpha 1$  subunit of GABA<sub>A</sub> receptors. *J Neurosci* **26**:5673–5683.
- Pritchett DB, Sontheimer H, Shivers BD, Ymer S, Kettenmann H, Schofield PR, and Seeburg PH (1989) Importance of a novel GABA<sub>A</sub> receptor subunit for benzodiazepine pharmacology. *Nature* **338**:582–585.
- Quinlan JJ, Firestone LL, and Homanics GE (2000) Mice lacking the long splice variant of the  $\gamma 2$  subunit of the GABA<sub>A</sub> receptor are more sensitive to benzodiazepines. *Pharmacol Biochem Behav* **66**:371–374.
- Quirk K, Gillard NP, Ragan CI, Whiting PJ, and McKernan RM (1994)  $\gamma$ -Aminobutyric acid type A receptors in the rat brain can contain both  $\gamma 2$  and  $\gamma 3$  subunits, but  $\gamma 1$  does not exist in combination with another  $\gamma$  subunit. *Mol Pharmacol* **45**:1061–1070.
- Reynolds DS, Rosahl TW, Cirone J, O'Meara GF, Haythornthwaite A, Newman RJ, Myers J, Sur C, Howell O, Rutter AR, et al. (2003) Sedation and anesthesia mediated by distinct GABA<sub>A</sub> receptor isoforms. *J Neurosci* **23**:8608–8617.
- Rhodes KJ and Trimmer KS (2006) Antibodies as valuable neuroscience research tools versus reagents of mass distraction. *J Neurosci* **26**:8017–8020.
- Rivera C, Voipio J, and Kaila K (2005) Two developmental switches in GABAergic signalling: the K<sup>+</sup>-Cl<sup>-</sup> cotransporter KCC2 and carbonic anhydrase CAVII. *J Physiol* **562**:27–36.
- Robinson TN and Olsen RW (1988) GABA, in *Comparative Invertebrate Neurochemistry* (Lunt GG, and Olsen RW eds) pp 90–123, Croom Helm, London.
- Rudolph U and Antkowiak B (2004) Molecular and neuronal substrates for general anaesthetics. *Nat Rev Neurosci* **5**:709–720.
- Rudolph U, Crestani F, Benke D, Brünig I, Benson JA, Fritschy JM, Martin JR, Blüthmann H, and Möhler H (1999) Benzodiazepine actions mediated by specific GABA<sub>A</sub> receptor subtypes. *Nature* **401**:796–800.
- Rudolph U and Möhler H (2004) Analysis of GABA<sub>A</sub> receptor function, and dissection of the pharmacology of benzodiazepines and general anesthetics through mouse genetics. *Annu Rev Pharmacol Toxicol* **44**:475–498.
- Saxena NC and Macdonald RL (1994) Assembly of GABA<sub>A</sub> receptor subunits: role of  $\delta$  subunit. *J Neurosci* **14**:7077–7086.
- Semyanov A, Walker MC, Kullmann DM, and Silver RA (2004) Tonically active GABA<sub>A</sub> receptors: modulating gain and maintaining the tone. *Trends Neurosci* **27**:262–269.
- Sergeeva OA, Andreeva N, Garret M, Scherer A, and Haas HL (2005) Pharmacological properties of GABA<sub>A</sub> receptors in rat hypothalamic neurons expressing the  $\epsilon$  subunit. *J Neurosci* **25**:88–95.
- Sieghart W (1989) Multiplicity of GABA<sub>A</sub> benzodiazepine receptors. *Trends Pharmacol Sci* **10**:407–411.
- Sieghart W (1995) Structure and pharmacology of GABA<sub>A</sub> receptor subtypes. *Pharmacol Rev* **47**:181–234.
- Sieghart W and Ernst M (2005) Heterogeneity of GABA<sub>A</sub> receptors: revived interest in the development of subtype-selective drugs. *Curr Med Chem Cent Nerv Syst Agents* **5**:217–242.

- Sieghart W and Sperk G (2002) Subunit composition, distribution and function of GABA<sub>A</sub> receptors. *Curr Top Med Chem* **2**:795–816.
- Sigel E (2002) Mapping of the benzodiazepine recognition site on GABA<sub>A</sub> receptor. *Curr Top Med Chem* **2**:833–839.
- Sigel E and Buhr A (1997) The benzodiazepine binding site of GABA<sub>A</sub> receptors. *Trends Pharmacol Sci* **18**:425–429.
- Sigel E, Baur R, Boulineau N, and Minier F (2006) Impact of subunit positioning on GABA<sub>A</sub> receptor function. *Biochem Soc Trans* **34**:868–871.
- Sigel E, Baur R, Kellenberger S, and Malherbe P (1992) Point mutations affecting antagonist and agonist dependent gating of the GABA<sub>A</sub> receptor channels. *EMBO J* **11**:2017–2023.
- Simon J, Wakimoto H, Fujita N, Lalande M, and Barnard EA (2004) Analysis of the set of GABA<sub>A</sub> receptor genes in the human genome. *J Biol Chem* **279**:41422–41435.
- Sine S and Engel A (2006) Recent advances in Cys-loop receptor structure and function. *Nature* **440**:448–455.
- Sinkkonen ST, Hanna MC, Kirkness EF, and Korpi ER (2000) GABA<sub>A</sub> receptor  $\epsilon$  and  $\theta$  subunits display unusual structural variations between species and are enriched in the rat locus ceruleus. *J Neurosci* **20**:3588–3595.
- Smith AJ, Oxley B, Malpas S, Pillai GV, and Simpson PB (2004) Compounds exhibiting selective efficacy for different  $\beta$  subunits of human recombinant GABA<sub>A</sub> receptors. *J Pharmacol Exp Ther* **311**:601–609.
- Smith GB and Olsen RW (1995) Functional domains of GABA<sub>A</sub> receptors. *Trends Pharmacol Sci* **16**:162–168.
- Sohal VS, Keist R, Rudolph U, and Huguenard JR (2003) Dynamic GABA<sub>A</sub> receptor subtype-specific modulation of the synchrony and duration of thalamic oscillations. *J Neurosci* **23**:3649–3657.
- Somogyi P, Fritschy JM, Benke D, Roberts JDB, and Sieghart W (1996) The  $\gamma$ 2 subunit of the GABA<sub>A</sub> receptor is concentrated in synaptic junctions containing the  $\alpha$ 1 and  $\beta$ 2/3 subunits in hippocampus, cerebellum and globus pallidus. *Neuropharmacology* **35**:1425–1444.
- Stell B, Brickley SG, Tang CY, Farrant M, and Mody I (2003) Neuroactive steroids reduce neuronal excitability by selectively enhancing tonic inhibition mediated by  $\delta$  subunit-containing GABA<sub>A</sub> receptors. *Proc Natl Acad Sci U S A* **100**:14439–14444.
- Stephens DN, Pistovcakova J, Worthing L, Atack JR, and Dawson GR (2006) Role of GABA<sub>A</sub>  $\alpha$ 5-containing receptors in ethanol reward: the effects of targeted gene deletion, and a selective inverse agonist. *Eur J Pharmacol* **526**:240–250.
- Sur C, Wafford KA, Reynolds DS, Hadingham KL, Bromidge F, Macaulay A, Collinson N, O'Meara G, Howell O, Newman R, et al. (2001) Loss of the major GABA<sub>A</sub> receptor subtype in the brain is not lethal in mice. *J Neurosci* **21**:3409–3418.
- Sur C, Wuirk K, Dewar D, Atack J, and McKernan R (1998) Rat and human hippocampal  $\alpha$ 5 subunit-containing GABA<sub>A</sub> receptors have  $\alpha$ 5 $\beta$ 3 $\gamma$ 2 pharmacological characteristics. *Mol Pharmacol* **54**:928–933.
- Thompson AJ and Lummis SCR (2006) 5-HT<sub>3</sub> receptors. *Curr Pharm Des* **12**:3615–3630.
- Thompson SA, Smith MZ, Wingrove PB, Whiting PJ, and Wafford KA (1999) Mutation at the putative GABA<sub>A</sub> ion-channel gate reveals changes in allosteric modulation. *Br J Pharmacol* **127**:1349–1358.
- Thompson SA, Wheat L, Brown NA, Wingrove PB, Pillai GV, Whiting PJ, Adkins C, Woodward CH, Smith AJ, Simpson PB, Collins I, and Wafford KA (2004) Salicylidene salicylhydrazide, a selective inhibitor of  $\beta$ 1-containing GABA<sub>A</sub> receptors. *Br J Pharmacol* **142**:97–106.
- Thompson SA, Whiting PJ, and Wafford KA (1996) Barbiturate interactions at the human GABA<sub>A</sub> receptor: dependence on receptor subunit combination. *Br J Pharmacol* **117**:521–527.
- Tierney ML, Birnir B, Pillai NP, Clements JD, Howitt SM, Cox GB, and Gage PW (1996) Effects of mutating leucine to threonine in the M2 segment of  $\alpha$ 1 and  $\beta$ 1 subunits of GABA<sub>A</sub>  $\alpha$ 1 $\beta$ 1 receptors. *J Membr Biol* **154**:11–21.
- Tögel M, Mossier B, Fuchs K, and Sieghart W (1994)  $\gamma$ -Aminobutyric acid receptors displaying association of  $\gamma$ 3-subunits with  $\beta$ 2/3- and different  $\alpha$ -subunits exhibit unique pharmacological properties. *J Biol Chem* **269**:12993–12998.
- Tretter V, Hauer B, Nusser Z, Mihalek RM, Höger H, Homanics GE, Somogyi P, and Sieghart W (2001) Targeted disruption of the GABA<sub>A</sub> receptor  $\delta$  subunit gene leads to an up-regulation of  $\gamma$ 2 subunit-containing receptors in cerebellar granule cells. *J Biol Chem* **276**:10532–10538.
- Unwin N (2005) Refined structure of the nicotinic acetylcholine receptor at 4 Å resolution. *J Mol Biol* **346**:967–989.
- Vicini S, Ferguson C, Prybylowski K, Kralic J, Morrow AL, and Homanics GE (2001) GABA<sub>A</sub> receptor  $\alpha$ 1 subunit deletion prevents developmental changes of inhibitory synaptic currents in cerebellar neurons. *J Neurosci* **21**:3009–3016.
- Wagner DA, Goldschen-Ohm MP, Hales TG, and Jones MV (2005) Kinetics and spontaneous open probability conferred by the epsilon subunit of the GABA<sub>A</sub> receptor. *J Neurosci* **25**:10462–10468.
- Wallner M, Hanchar H, and Olsen RW (2003) Ethanol enhances  $\alpha$ 4 $\beta$ 3 $\delta$  and  $\alpha$ 6 $\beta$ 3 $\delta$  GABA<sub>A</sub> receptors at low concentrations known to affect humans. *Proc Natl Acad Sci U S A* **100**:15218–15223.
- Wallner M, Hanchar HJ, and Olsen RW (2006) Low dose alcohol actions on  $\alpha$ 4 $\beta$ 3 $\delta$  GABA<sub>A</sub> receptors are reversed by the behavioral alcohol antagonist Ro15-4513. *Proc Natl Acad Sci U S A* **103**:8540–8545.
- Wei W, Zhang N, Peng Z, Houser CR, and Mody I (2003) Perisynaptic localization of  $\delta$  subunit-containing GABA<sub>A</sub> receptors and their activation by GABA spillover in the mouse dentate gyrus. *J Neurosci* **23**:10650–10661.
- Whiting PJ (2006) GABA<sub>A</sub> receptors: a viable target for novel anxiolytics? *Curr Opin Pharmacol* **6**:24–29.
- Whiting PJ, McAllister G, Vassilatis D, Bonnert TP, Heavens RP, Smith DW, Hewson L, O'Donnell R, Rigby MR, Sirinathsinghji DJ, et al. (1997) Neuronally restricted RNA splicing regulates the expression of a novel GABA<sub>A</sub> receptor subunit conferring atypical functional properties. *J Neurosci* **17**:5027–5037.
- Whiting PJ, McKernan RM, and Iversen LL (1990) Another mechanism for creating diversity in  $\gamma$ -aminobutyrate type A receptors: RNA splicing directs expression of two forms of  $\gamma$ 2 phosphorylation site. *Proc Natl Acad Sci U S A* **87**:330–337.
- Whiting PJ, Wafford KA, and McKernan RM (2000) Pharmacological subtypes of GABA<sub>A</sub> receptors based on subunit composition, in *GABA in the Nervous System: The View at Fifty Years* (Martin DL, Olsen RW, eds) pp. 113–126, Lippincott Williams & Wilkins, Philadelphia.
- Wingrove PB, Wafford KA, and Bain C (1994) The modulatory action of loreclezole at the GABA<sub>A</sub> receptor is determined by a single amino acid in the  $\beta$ 2 and  $\beta$ 3 subunits. *Proc Natl Acad Sci U S A* **91**:4569–4573.
- Wisden W, Laurie DJ, Monyer H, and Seeburg PH (1992) The distribution of 13 GABA<sub>A</sub> receptor subunit mRNAs in the rat brain: I. Telencephalon, diencephalon, mesencephalon. *J Neurosci* **12**:1040–1062.
- Wohlfarth KM, Bianchi MT, and Macdonald RL (2002) Enhanced neurosteroid potentiation of ternary GABA<sub>A</sub> receptors containing the  $\delta$  subunit. *J Neurosci* **22**:1541–1549.
- Woodward RM, Polenzani L, and Miledi R (1992) Effects of steroids on  $\gamma$ -aminobutyric acid receptors expressed in *Xenopus* oocytes by poly<sup>A</sup> RNA from mammalian brain and retina. *Mol Pharmacol* **41**:89–103.
- Yee BK, Keist R, von Böhrer L, Studer R, Benke D, Hagenbuch N, Dong Y, Malenka RC, Fritschy JM, Blüthmann H, et al. (2005) A schizophrenia-related sensorimotor deficit links  $\alpha$ 3-containing GABA<sub>A</sub> receptors to a dopamine hyperfunction. *Proc Natl Acad Sci U S A* **102**:17154–17159.
- Zhang JH, Sato M, and Tohyama M (1991) Different postnatal development profiles of neurons containing distinct GABA<sub>A</sub> receptor  $\beta$  subunit mRNAs ( $\beta$ 1,  $\beta$ 2, and  $\beta$ 3) in the rat forebrain. *J Comp Neurol* **308**:586–613.